

# MMPI-2-RF® Bibliography

## MMPI-2-RF REFERENCES BY TOPIC

### BASIC SOURCES

- Ben-Porath, Y. S. (2012). *Interpreting the MMPI-2-RF*. Minneapolis, MN: University of Minnesota Press.
- Ben-Porath, Y. S., & Tellegen, A. (2008/2011). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2-Restructured Form): Manual for administration, scoring, and interpretation*. Minneapolis, MN: University of Minnesota Press.
- Ben-Porath, Y. S., & Tellegen, A. (2011). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2-Restructured Form): User's guide for reports* (2nd ed.). Minneapolis, MN: University of Minnesota Press.
- Block, A. R., & Ben-Porath, Y. S. (2018). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2-Restructured Form): User's guide for the Spine Surgery and Spinal Cord Stimulator Candidate Interpretive Reports*. Minneapolis, MN: University of Minnesota Press.
- Corey, D. M., & Ben-Porath, Y. S. (2014). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2-Restructured Form): User's guide for the Police Candidate Interpretive Report*. Minneapolis, MN: University of Minnesota Press.
- Tellegen, A., & Ben-Porath, Y. S. (2008/2011). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2-Restructured Form): Technical manual*. Minneapolis, MN: University of Minnesota Press.
- Tellegen, A., Ben-Porath, Y. S., McNulty, J. L., Arbisi, P. A., Graham, J. R., & Kaemmer, B. (2003). *MMPI-2 Restructured Clinical (RC) Scales: Development, validation, and interpretation*. Minneapolis, MN: University of Minnesota Press.

### BOOKS

- Ben-Porath, Y. S. (2012). *Interpreting the MMPI-2-RF*. Minneapolis, MN: University of Minnesota Press.
- Corey, D. M., & Ben-Porath, Y. S. (2018). *Assessing police and other public safety personnel using the MMPI-2-RF*. Minneapolis, MN: University of Minnesota Press.
- Friedman, A. F., Bolinsky, P. K., Levak, R., & Nichols, D. S. (2015). *Psychological assessment with the MMPI-2/RF (Third Edition)*. New York, NY: Routledge/Taylor & Francis.
- Graham, J. R. (2012). *MMPI-2: Assessing personality and psychopathology* (5th ed.). New York, NY: Oxford University Press.
- Greene, R. L. (2011). *MMPI-2/MMPI-2-RF: An interpretive manual* (3<sup>rd</sup> ed.). New York, NY: Allyn & Bacon.

- McCord, D. M. (2018). *Assessment Using the MMPI-2-RF*. Washington, DC: American Psychological Association.
- Sellbom, M., & Wygant, D. B. (2018). *Forensic applications of the MMPI-2-RF: A casebook*. Minneapolis, MN: University of Minnesota Press.

## BOOK CHAPTERS

- Ben-Porath, Y. S. (2013). Forensic applications of the Minnesota Multiphasic Personality Inventory-2-Restructured Form. In R. P. Archer and E. M. A. Wheeler (Eds.), *Forensic use of clinical assessment instruments* (pp. 63–107). New York, NY: Routledge.
- Ben-Porath, Y. S. (2013). The MMPI instruments. In S. Koffler, J. Morgan, I. S. Barron, & M. F. Greiffenstein (Eds.), *Neuropsychology: Science and practice, I*. New York, NY: Oxford University Press.
- Ben-Porath, Y. S. (2013). Understanding and using the MMPI-2-RF. In G. P. Koocher, J. C. Norcross, & B. A. Greene (Eds.), *Psychologists' desk reference* (pp. 12–133). New York, NY: Oxford University Press.
- Ben-Porath, Y. S., & Archer, R. P. (2014). The MMPI instruments. In R.P. Archer and S.R. Smith (eds.) *Personality Assessment*, 2nd edition (pp. 89-146). NY: Routledge.
- Ben-Porath, Y. S., Corey, D. M., & Tarescavage, A. M. (2017). Using the MMPI-2-RF in preemployment evaluations of police officer candidates. In C. L. Mitchell & E. H. Dorian (eds.) *Police psychology and its growing impact on modern law enforcement*, (pp. 51-78). Hershey, PA: IGI Global. doi: [10.4018/978-1-5225-0813-7.ch003](https://doi.org/10.4018/978-1-5225-0813-7.ch003)
- Crighton, A. C., & Ben-Porath, Y. S. (2016). The MMPI-2-RF Substantive Scales. In *Encyclopedia of Personality and Individual Differences*. (pp. 1-5). New York, NY: Springer International Publishing.
- Gervais, R. O., Louw, D., & Gibson, K. (2017). Using the MMPI-2-RF in discriminating between malingering and somatoform disorder. In K. B. Boone (Ed.) *Neuropsychological Evaluation of Somatoform and Other Functional Somatic Conditions*. NY: Routledge.
- Marek, R. J., & Ben-Porath, Y. S. (2017). Using the MMPI-2 Restructured Form (MMPI-2-RF) in behavioral medicine setting. In M. Maurush (Ed.) *Handbook of Psychological Assessment in Primary Care Settings* (2nd edition, pp. 631-661). NY: Taylor and Francis.
- McCord, D. M. (2013). Minnesota Multiphasic Personality Inventory. In D. S. Dunn (Ed.), *Oxford Bibliographies in Psychology*. New York, NY: Oxford University Press. doi: [10.1093/OBO/9780199828340-0118](https://doi.org/10.1093/OBO/9780199828340-0118)
- Sellbom, M., & Lee, T. T. C. (2013). Assessment of anxiety symptoms using the MMPI-2, MMPI-2-RF, and MMPI-A. In D. McKay & E. A. Storch (Eds.), *Handbook of assessing variants and complications in anxiety disorders* (pp. 139–162). New York, NY: Springer.
- Tarescavage, A. M., & Ben-Porath, Y. S. (2014). Minnesota Multiphasic Personality Inventory instruments. In R. L. Cautin & S. O. Lillienfeld (Eds.). *The encyclopedia of clinical psychology*. Wiley Online Library. doi: [10.1002/9781118625392.wbecp265](https://doi.org/10.1002/9781118625392.wbecp265)
- Wygant, D. B., Walls, B. D., Brothers, S. L., & Berry, D. T. R. (2018). Assessment of Malingering and Defensiveness on the MMPI-2 and MMPI-2-RF. R. Rogers & S. Bender (Eds.), *Clinical Assessment of Malingering and Deception*, 4th ed. (p. 257-279). New York: Guilford Press.

## PEER-REVIEWED PUBLICATIONS

### GENERAL ISSUES

- Anderson, J. L., Sellbom, M., Bagby, R. M., Quilty, L. C., Veltri, C. O. C., Markon, K. E., & Krueger, R. F. (2013). On the convergence between PSY-5 domains and PID-5 domains and facets: Implications for assessment of DSM-5 personality traits. *Assessment, 20*, 286–294. doi: [10.1177/1073191112471141](https://doi.org/10.1177/1073191112471141)
- Arbisi, P. A. (2014). Introduction to the special section on the Personality Psychopathology Five (PSY-5) and DSM-5 trait dimensional diagnostic systems for personality disorders: Emerging convergence. *Journal of Personality Assessment, 96*, 117–120. doi: [10.1080/00223891.2013.866571](https://doi.org/10.1080/00223891.2013.866571)
- Archer, R. P. (2006). A perspective on the Restructured Clinical (RC) Scale project. *Journal of Personality Assessment, 87*, 179–185. doi: [10.1207/s15327752jpa8702\\_07](https://doi.org/10.1207/s15327752jpa8702_07)
- Ben-Porath, Y. S. (2017). An update to Williams and Lally's (2017) analysis of MMPI-2-RF acceptance. *Professional Psychology Research and Practice, 48*, 275–278. doi: [10.1037/pro0000115](https://doi.org/10.1037/pro0000115)
- Ben-Porath, Y. S., & Tellegen, A. (2008). Empirical correlates of the MMPI-Restructured Clinical (RC) Scales in mental health, forensic, and non-clinical settings: An introduction. *Journal of Personality Assessment, 90*, 119–121. doi: [10.1080/00223890701845120](https://doi.org/10.1080/00223890701845120)
- Bolinger, E., Reese, C., Suhr, J., & Larrabee, G. J. (2014). Susceptibility of the MMPI-2-RF Neurological Complaints and Cognitive Complaints Scales to over-reporting in simulated head injury. *Archives of Clinical Neuropsychology, 29*, 7–15. doi: [10.1093/arclin/act082](https://doi.org/10.1093/arclin/act082)
- Bolinski, P. K., & Nichols, D. S. (2011). Construct drift in the MMPI-2 Restructured Clinical Scales: Further evidence and a possible historic example. *Journal of Clinical Psychology, 67*, 1–11. doi: [10.1002/jclp.20814](https://doi.org/10.1002/jclp.20814)
- Burchett, D. L., & Ben-Porath, Y. S. (2010). The impact of overreporting on MMPI-2-RF substantive scale score validity. *Assessment, 17*, 497–516. doi: [10.1177/1073191110378972](https://doi.org/10.1177/1073191110378972)
- Caldwell, A. B. (2006). Maximal measurement or meaningful measurement: The interpretive challenges of the MMPI-2 Restructured Clinical (RC) Scales. *Journal of Personality Assessment, 87*, 193–201. doi: [10.1207/s15327752jpa8702\\_09](https://doi.org/10.1207/s15327752jpa8702_09)
- Chmielewski, M., Bagby, R. M., Markon, K., Ring, A. J., & Ryder, A. G. (2014). Openness to experience, intellect, schizotypal personality disorder, and psychoticism: Resolving the controversy. *Journal of Personality Disorders, 28*, 128. doi: [10.1521/pedi\\_2014\\_28\\_128](https://doi.org/10.1521/pedi_2014_28_128)
- Cox, A., Courge, S. C., Felder, A. H., & Weed, N. C. (2017). Effects of augmenting response options of the MMPI-2-RF: An extension of previous findings. *Cogent Psychology, 4*, 1323988. doi: [10.1080/23311908.2017.1323988](https://doi.org/10.1080/23311908.2017.1323988)
- Cox, A. Pant, H., Gilson, A. N., Rodriguez, J. S., Young, K. R., Kwon, S., & Weed, N. C., (2012). Effects of augmenting response options on MMPI-2 RC Scale psychometrics. *Journal of Personality Assessment, 94*, 613–619. doi: [10.1080/00223891.2012.700464](https://doi.org/10.1080/00223891.2012.700464)
- Eaton, N., Krueger, R. F., South, S. C., & Simms, L. J. (2011). Contrasting prototypes and dimensions in the classification of personality pathology: Evidence that dimensions, but not prototypes, are robust. *Psychological Medicine, 41*, 1151–1163. doi: [10.1017/S0033291710001650](https://doi.org/10.1017/S0033291710001650)

- Finn, J. A., Ben-Porath, Y. S., & Tellegen, A. (2015). Dichotomous versus polytomous response options in psychopathology assessment: Method or meaningful variance? *Psychological Assessment, 27*, 184–193. doi: [10.1037/pas0000044](https://doi.org/10.1037/pas0000044)
- Finn, S. E., & Kamphuis, J. H. (2006). The MMPI-2 RC Scales and restraints to innovation, or “What have they done to my song?” *Journal of Personality Assessment, 87*, 202–210. doi: [10.1207/s15327752jpa8702\\_10](https://doi.org/10.1207/s15327752jpa8702_10)
- Forbey, J. D., & Ben-Porath, Y. S. (2007). Computerized adaptive personality testing: A review and illustration with the MMPI-2 Computerized Adaptive Version (MMPI-2-CA). *Psychological Assessment, 19*, 14–24. doi: [10.1037/1040-3590.19.1.14](https://doi.org/10.1037/1040-3590.19.1.14)
- Greene, R. L., Rouse, S. V., Butcher, J. N., Nichols, D. S., & Williams, C. L. (2009). The MMPI-2 Restructured Clinical (RC) Scales and redundancy: Response to Tellegen, Ben-Porath, and Sellbom. *Journal of Personality Assessment, 91*, 222–226. doi: [10.1080/00223890902800825](https://doi.org/10.1080/00223890902800825)
- Harkness, A. R., Finn, J. A., McNulty, J. L., & Shields, S. M. (2012). The Personality Psychopathology Five (PSY-5): Recent constructive replication and assessment literature review. *Psychological Assessment, 24*, 432–443. doi: [10.1037/a0025830](https://doi.org/10.1037/a0025830)
- Harkness, A. R., McNulty, J. L., Finn, J. A., Reynolds, S. M., & Shields, S. M. (2014). The MMPI-2-RF Personality Psychopathology Five (PSY-5) Scales: Development and validity research. *Journal of Personality Assessment, 96*, 140–150. doi: [10.1080/00223891.2013.823439](https://doi.org/10.1080/00223891.2013.823439)
- Harkness, A. R., Reynolds, S. M., & Lilienfeld, S. O. (2014). A review of systems for psychology and psychiatry: Adaptive systems, Personality Psychopathology-Five (PSY-5), and DSM-5. *Journal of Personality Assessment, 96*, 121–139. doi: [10.1080/00223891.2013.823438](https://doi.org/10.1080/00223891.2013.823438)
- Hoelzle, J. B., & Meyer, G. J. (2008). The factor structure of the MMPI-2 Restructured Clinical (RC) Scales. *Journal of Personality Assessment, 90*, 443–455. doi: [10.1080/00223890802248711](https://doi.org/10.1080/00223890802248711)
- Lally, S. J., & Williams, C. L. (2017). Response to Ben-Porath’s update to Williams and Lally (2017). *Professional Psychology Research and Practice, 48*, 282–285. doi: [10.1037/pro0000157](https://doi.org/10.1037/pro0000157)
- Lanyon, R. I., & Thomas, M. I. (2013). Assessment of global psychiatric categories: The PSI/PSI-2 and the MMPI-2-RF. *Psychological Assessment, 25*, 227–232. doi: [10.1037/a0030313](https://doi.org/10.1037/a0030313)
- Lee, T. C., Forbey, J. D., & Ritchey, K. A. (2011). The impact of emotional priming on MMPI-2 scale scores. *Assessment, 18*, 102–105. doi: [10.1177/1073191110392496](https://doi.org/10.1177/1073191110392496)
- Martin, P. K., Schroeder, R. W., & Odland, A. P. (2015). Neuropsychologists’ validity testing beliefs and practices: A survey of North American professionals. *The Clinical Neuropsychologist, 29*, 741–776. doi: [10.1080/13854046.2015.1087597](https://doi.org/10.1080/13854046.2015.1087597)
- Menton, W. H., Crighton, A. H., Tarescavage, A. M., Marek, R. J., Hicks, A. D., & Ben-Porath, Y. S. (2017). Equivalence of laptop and tablet administrations of the Minnesota Multiphasic Personality Inventory-2 Restructured Form. *Assessment*. doi: [0.1177/1073191117714558](https://doi.org/10.1177/1073191117714558)
- McNulty, J. L., & Overstreet, S. R. (2014). Viewing the MMPI-2-RF structure through the Personality Psychopathology Five (PSY-5) lens. *Journal of Personality Assessment, 96*, 151–157. doi: [10.1080/00223891.2013.840305](https://doi.org/10.1080/00223891.2013.840305)
- Nichols, D. S. (2006a). The trials of separating bath water from baby: A review and critique of the MMPI-2 Restructured Clinical Scales. *Journal of Personality Assessment, 87*, 121–138. doi: [10.1207/s15327752jpa8702\\_02](https://doi.org/10.1207/s15327752jpa8702_02)

- Nichols, D. S. (2006b). Commentary on Rogers, Sewell, Harrison, & Jordan (2006). *Journal of Personality Assessment*, *87*, 172–174. doi: [10.1207/s15327752jpa8702\\_05](https://doi.org/10.1207/s15327752jpa8702_05)
- Odland, A. P., Lammy, A. B., Perle, J. G., Martin, P. K., & Grote, C. L. (2015). Reaffirming normal: The high risk of pathologizing healthy adults when interpreting the MMPI-2-RF. *The Clinical Neuropsychologist*. doi: [10.1080/13854046.2015.1005675](https://doi.org/10.1080/13854046.2015.1005675)
- Patrick, R. E., & Horner, M. D. (2014). Psychological characteristics of individuals who put forth inadequate effort in a secondary gain context. *Archives of Clinical Neuropsychology*, *29*, 754–756. doi: [10.1093/arclin/acu054](https://doi.org/10.1093/arclin/acu054)
- Rogers, R., & Sewell, K. W. (2006). MMPI-2 at the crossroads: Aging technology or radical retrofitting. *Journal of Personality Assessment*, *87*, 175–178. doi: [10.1207/s15327752jpa8702\\_06](https://doi.org/10.1207/s15327752jpa8702_06)
- Rogers, R., Sewell, K. W., Harrison, K. S., & Jordan, M. J. (2006). The MMPI-2 Restructured Clinical Scales: A paradigmatic shift in scale development. *Journal of Personality Assessment*, *87*, 139–147. doi: [10.1207/s15327752jpa8702\\_03](https://doi.org/10.1207/s15327752jpa8702_03)
- Rouse, S. V., Greene, R. L., Butcher, J. N., Nichols, D. S., & Williams, C. L. (2008). What do the MMPI-2 Restructured Clinical Scales reliably measure? Answers from multiple research settings. *Journal of Personality Assessment*, *90*, 435–442. doi: [10.1080/00223890802248695](https://doi.org/10.1080/00223890802248695)
- Sellbom, M., Anderson, J. L., & Bagby, R. M. (2013). Assessing DSM-5 Section III personality traits and disorders with the MMPI-2-RF. *Assessment*, *20*, 709–722. doi: [10.1177/1073191113508808](https://doi.org/10.1177/1073191113508808)
- Sellbom, M., & Arbisi, P. A. (in press). Introduction to the special issue: Linking the MMPI-2-RF to contemporary models of psychopathology. *Journal of Personality Assessment*.
- Sellbom, M., Ben-Porath, Y. S. & Bagby, R. M. (2008). On the hierarchical structure of mood and anxiety disorders: Confirmatory evidence and elaboration of a model of temperament markers. *Journal of Abnormal Psychology*, *117*, 576–590. doi: [10.1037/a0012536](https://doi.org/10.1037/a0012536)
- Sellbom, M., Ben-Porath, Y. S., Graham, J. R., Arbisi, P. A., & Bagby, R. M. (2005). Susceptibility of the MMPI-2 Clinical, Restructured Clinical (RC), and Content Scales to overreporting and underreporting. *Assessment*, *12*, 79–85. doi: [10.1177/1073191104273515](https://doi.org/10.1177/1073191104273515)
- Simms, L. J. (2006). Bridging the divide: Comments on the Restructured Clinical Scales of the MMPI-2. *Journal of Personality Assessment* *87*, 211–216. doi: [10.1207/s15327752jpa8702\\_11](https://doi.org/10.1207/s15327752jpa8702_11)
- Tarescavage, A. M. & Ben-Porath, Y. S. (2015). A response to Odland et al.'s misleading, alarmist estimates of risk for overpathologizing when interpreting the MMPI-2-RF. *The Clinical Neuropsychologist*, *29*, 183–196. doi: [10.1080/13854046.2015.1040843](https://doi.org/10.1080/13854046.2015.1040843)
- Tarescavage, A. M. & Ben-Porath, Y. S. (2017). Examination of the feasibility and utility of Flexible and Conditional Administration (FCA) of the Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF). *Psychological Assessment*, *29*, 1337-1348. doi: [10.1037/pas0000442](https://doi.org/10.1037/pas0000442)
- Tarescavage, A. M., Marek, R. J., Finn, J. A., Hicks, A., Rapier, J. L., & Ben-Porath, Y. S. (2013). Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) normative elevation rates: Comparisons with epidemiological prevalence rates. *The Clinical Neuropsychologist*, *27*, 1106–1120. doi: [10.1080/13854046.2013.832386](https://doi.org/10.1080/13854046.2013.832386)
- Tellegen, A., Ben-Porath, Y. S., & Sellbom, M. (2009). Construct validity of the MMPI-2 Restructured Clinical (RC) Scales: Reply to Rouse, Greene, Butcher, Nichols, & Williams. *Journal of Personality Assessment*, *91*, 211–221. doi: [10.1080/00223890902794192](https://doi.org/10.1080/00223890902794192)



- Tellegen, A., Ben-Porath, Y. S., Sellbom, M., Arbisi, P. A., McNulty, J. L., & Graham, J. R. (2006). Further evidence on the validity of the MMPI-2 Restructured Clinical (RC) Scales: Addressing questions raised by Rogers et al. and Nichols. *Journal of Personality Assessment, 87*, 148–171. doi: [10.1207/s15327752jpa8702\\_04](https://doi.org/10.1207/s15327752jpa8702_04)
- Trumbetta, S. L., Bolinsky, P. K., & Gottesman, I. I. (2013). The MMPI-ARC and MMPI-AF Scales: Psychometric properties of the MMPI-2-RF's Restructured Clinical (RC) and Higher-Order Factor Scales when adapted for use with archival adolescent MMPI data. *Archives of Assessment Psychology, 3*, 24–36.
- Van der Heijden, P. T., Egger, J. I. M., & Derksen, J. J. L. (2008). Psychometric evaluation of the MMPI-2 Restructured Clinical Scales in two Dutch samples. *Journal of Personality Assessment, 90*, 456–464. doi: [10.1080/00223890802248745](https://doi.org/10.1080/00223890802248745)
- Van der Heijden, P. T., Egger, J. I. M., & Derksen, J. J. L. (2010). Comparability of scores on the MMPI-2-RF scales generated with the MMPI-2 and MMPI-2-RF booklets. *Journal of Personality Assessment, 92*, 254–259. doi: [10.1080/00223891003670208](https://doi.org/10.1080/00223891003670208)
- Van der Heijden, P. T., Rossi, G. M., Van der Veld, M. M., Derksen, J. J. L., & Egger, J. I. M. (2013). Personality and psychopathology: Higher order relations between the Five Factor Model of personality and the MMPI-2 Restructured Form. *Journal of Research in Personality, 47*, 572–579. doi: [10.1016/j.jrp.2013.05.001](https://doi.org/10.1016/j.jrp.2013.05.001)
- Van der Heijden, P. T., Rossi, G. M., Van der Veld, M. M., Derksen, J. J. L., & Egger, J. I. M. (2013). Personality and psychopathology: Mapping the MMPI-2-RF on Cloninger's psychobiological model of personality. *Assessment, 20*, 576–584. doi: [10.1177/1073191113490791](https://doi.org/10.1177/1073191113490791)
- Viken, R. J., & Rose, R. J. (2007). Genetic variation and covariation in the original and Restructured Clinical Scales of the MMPI. *Journal of Abnormal Psychology, 116*, 842–847. doi: [10.1037/0021-843X.116.4.842](https://doi.org/10.1037/0021-843X.116.4.842)
- Weed, N. C. (2006). Syndromal complexity, paradigm shifts, and the future of validation research: A comment on Nichols and Rogers, Sewell, Harrison, & Jordan. *Journal of Personality Assessment, 87*, 217–222. doi: [10.1207/s15327752jpa8702\\_12](https://doi.org/10.1207/s15327752jpa8702_12)
- Weiss, P. A., Bell, K. J., & Weiss, W. U. (2010). Use of the MMPI-2 RC Scales to detect criminal malingering. *Journal of Police and Criminal Psychology, 25*, 49–55. doi: [10.1007/s11896-009-9056-9](https://doi.org/10.1007/s11896-009-9056-9)
- Williams, C. L., & Lally, S. J. (2017). MMPI-2, MMPI-2-RF, and MMPI-A Administrations (2007–2014): Any Evidence of a "New Standard?". *Professional Psychology: Research and Practice, 48*, 267–274. doi: [10.1037/pro0000088](https://doi.org/10.1037/pro0000088)

## VALIDITY SCALES

- Arbisi, P. A. (2017). Form vs Function, fighting the last war: a reflection on the exchange between Larrabee, Bianchini, Boone, and Rohling (2017) and Nichols (2017) over Nichols and Gass (2015). The Fake Bad Scale (FBS): malingering or ligation response syndrome – Which is it? *The Clinical Neuropsychologist, 31*, 1406–1411. doi: [10.1080/13854046.2017.1365933](https://doi.org/10.1080/13854046.2017.1365933)
- Armistead-Jehle, O., Grills, C. E., Bieu, R. K., & Kulas, J. F. (2016). Clinical utility of Memory Complaints Inventory to detect invalid test performance. *The Clinical Neuropsychologist, 30*, 610–628. doi: [10.1080/13854046.2016.1177597](https://doi.org/10.1080/13854046.2016.1177597)

- Armistead-Jehle, P., Cole, W.R., & Stegman, R.L. (2017). Performance and symptom validity testing as a function of Medical Board Evaluation in U.S. military service members with a history of mild traumatic brain injury. *Archives of Clinical Neuropsychology*, *33*, 120-124. doi: [10.1093/arclin/acx031](https://doi.org/10.1093/arclin/acx031)
- Armistead-Jehle, P., Cooper, D. B., Grills, C. E., Cole, W. R., Lippa, S. M., Stegman, R. L., & Lange, R. T. (2018). Clinical utility of the mBIAS and NSI validity-10 to detect symptom over-reporting following mild TBI: A multicenter investigation with military service members. *Journal of Clinical and Experimental Neuropsychology*, *40*, 213-223. doi: [10.1080/13803395.2017.1329406](https://doi.org/10.1080/13803395.2017.1329406)
- Armistead-Jehle, P., Grills, C. E., Bieu, R. K., & Kulas, J. F. (2016). Clinical utility of Memory Complaints Inventory to detect invalid test performance. *The Clinical Neuropsychologist*, *30*, 610-628. doi: [10.1080/13854046.2016.1177597](https://doi.org/10.1080/13854046.2016.1177597)
- Bianchini, K. J., Aguerrevere, L. E., Curtis, K. L., Roebuck-Spencer, T. M., Frey, F. C., Greve, K. W., & Calamia, M. (2017). Classification Accuracy of the Minnesota Multiphasic Personality Inventory-2 (MMPI-2)-Restructured Form Validity Scales in Detecting Malingered Pain-Related Disability. *Psychological Assessment*. doi: [10.1037/pas0000532](https://doi.org/10.1037/pas0000532)
- Blasco Sáiz, J. L., & Pallardó Durá, L. (2013). Detección de exageración de síntomas mediante el SIMS y el MMPI-2-RF en pacientes diagnosticados de trastorno mixto ansioso-depresivo y adaptativo en el contexto medicolegal: Un estudio preliminar [Symptom exaggeration detection by the SIMS and the MMPI-2-RF in patients diagnosed of mixed anxiety-depressive disorder and adjustment disorder in the medico-legal context: A preliminary study]. *Clínica y Salud*, *24*(3), 177-183. doi: [10.1016/S1130-5274\(13\)70019-7](https://doi.org/10.1016/S1130-5274(13)70019-7)
- Bodapati, A. S., Combs, H. L., Pastorek, N. J., Miller, B., Troyanskaya, M., Romesser, J., Sim, A., & Linck, J. (2018). Detection of symptom over-reporting on the Neurobehavioral Symptom Inventory in OEF/OIF/OND veterans with history of mild TBI. *The Clinical Neuropsychologist*. doi: [10.1080/13854046.2018.1482003](https://doi.org/10.1080/13854046.2018.1482003)
- Bridges, S. A., & Baum, L. J. (2013). An examination of the MMPI-2-RF L-r Scale in an outpatient protestant sample. *Journal of Psychology and Christianity*, *32*, 115-123.
- Burchett, D., Dragon, W. R., Smith Holbert, A. M., Tarescavage, A. M., Mattson, C. A., Handel, R. W., & Ben-Porath, Y. S. (2016). "False feigners": Examining the impact of non-content-based invalid responding on the Minnesota Multiphasic Personality Inventory-2 Restructured Form content-based invalid responding indicators. *Psychological Assessment*, *28*, 458-470. doi: [10.1037/pas0000205](https://doi.org/10.1037/pas0000205)
- Chmielewski, M., Zhu, J., Burchett, D., Bury, A.S., & Bagby, R. M. (2017). The comparative capacity of the Minnesota Multiphasic Personality Inventory-2 (MMPI-2) and MMPI-2 Restructured Form (MMPI-2-F) validity scales to detect malingering in a disability claimant sample. *Psychological Assessment*, *29*, 199-208. doi: [10.1037/pas0000328](https://doi.org/10.1037/pas0000328)
- Cook, C. M., Bolinger, E., & Suhr, J. (2016). Further validation of the Conner's Adult Attention Deficit/Hyperactivity Index (CAII) for detection of non-credible report of Attention Deficit/Hyperactivity Disorder symptoms. *Archives of Clinical Neuropsychology*, *31*, 358-364. doi: [10.1093/arclin/acw015](https://doi.org/10.1093/arclin/acw015)
- Crichton, A. H., Marek, R. J., Dragon, W. R., & Ben-Porath, Y. S. (2017). Utility of the MMPI-2-RF Validity Scales in the detection of simulated underreporting: Implications of incorporating a manipulation check. *Assessment*, *24*, 853-864. doi: [10.1177/107319111562701129](https://doi.org/10.1177/107319111562701129)

- Crichton, A. H., Tarescavage, A. M., Gervais, R. O., & Ben-Porath, Y. S. (2017). The generalizability of over-reporting across a battery of self-report measures: An investigation with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) and the Personality Assessment Inventory (PAI) in a civil disability sample. *Assessment, 24*, 555-574. doi: [10.1177/1073191115621791](https://doi.org/10.1177/1073191115621791)
- Dionysus, K., Denney, R. L., & Halfaker, D. A. (2011). Detecting negative response bias with the Fake Bad Scale, Response Bias Scale, Henry-Heilbronner Index of the Minnesota Multiphasic Personality Inventory-2. *Archives of Clinical Neuropsychology, 26*, 81-88. doi: [10.1093/arclin/acq](https://doi.org/10.1093/arclin/acq)
- Dragon, W. R., Ben-Porath, Y. S., & Handel, R. H. (2012). Examining the impact of unscorable item responses on the validity and interpretability of MMPI-2/MMPI-2-RF Restructured Clinical (RC) Scale scores. *Assessment, 19*, 101-113. doi: [10.1177/1073191111415362](https://doi.org/10.1177/1073191111415362)
- Forbey, J. D., Lee, T. T. C., Ben-Porath, Y. S., Arbisi, P. A., & Gartland, D. (2013). Associations between MMPI-2-RF Validity Scale scores and extra-test measures of personality and psychopathology. *Assessment, 20*, 448-461. doi: [10.1177/1073191113478154](https://doi.org/10.1177/1073191113478154)
- Gass, C. S., & Odland, A. P. (2012). Minnesota Multiphasic Personality Inventory-2 Revised Form [sic] Symptom Validity Scale—Revised (MMPI-2-RF FBS-r; also known as Fake Bad Scale): Psychometric characteristics in a nonlitigation neuropsychological setting. *Journal of Clinical and Experimental Neuropsychology, 34*, 561-570. doi: [10.1080/13803395.2012.666228](https://doi.org/10.1080/13803395.2012.666228)
- Gervais, R. O., Ben-Porath, Y. S., Wygant, D. B., & Green, P. (2007). Development and validation of a Response Bias Scale (RBS) for the MMPI-2. *Assessment, 14*, 196-208. doi: [10.1177/1073191106295861](https://doi.org/10.1177/1073191106295861)
- Gervais, R. O., Ben-Porath, Y. S., Wygant, D. B., & Green, P. (2008). Differential sensitivity of the Response Bias Scale and MMPI-2 Validity Scales to memory complaints. *The Clinical Neuropsychologist, 22*, 1061-1079. doi: [10.1080/13854040701756930](https://doi.org/10.1080/13854040701756930)
- Gervais, R. O., Ben-Porath, Y. S., & Wygant, D. B. (2009). Empirical correlates and interpretation of the MMPI-2-RF Cognitive Complaints (COG) Scale. *The Clinical Neuropsychologist, 23*, 996-1015. doi: [10.1080/13854040902748249](https://doi.org/10.1080/13854040902748249)
- Gervais, R. O., Ben-Porath, Y. S., Wygant, D. B., & Sellbom, M. (2010). Incremental validity of the MMPI-2-RF over-reporting scales and RBS in assessing the veracity of memory complaints. *Archives of Clinical Neuropsychology, 25*, 274-284. doi: [10.1093/arclin/acq018](https://doi.org/10.1093/arclin/acq018)
- Gervais, R. O., Tarescavage, A. M., Greiffenstein, M. F., Wygant, D. B., Deslauriers, C., & Arends, P. (2018). Inconsistent responding on the MMPI-2-RF and uncooperative attitude: Evidence from cognitive performance validity measures. *Psychological Assessment, 30*, 410-415. doi: [10.1037/pas0000506](https://doi.org/10.1037/pas0000506)
- Gervais, R. O., Wygant, D. B., Sellbom, M., & Ben-Porath, Y. S. (2011). Associations between Symptom Validity Test failure and scores on the MMPI-2-RF Validity and Substantive Scales. *Journal of Personality Assessment, 93*, 508-517. doi: [10.1080/00223891.2011.594132](https://doi.org/10.1080/00223891.2011.594132)
- Glassmire, D. M., Jhawar, A., Burchett, D., & Tarescavage, A. M. (2016). Evaluating item endorsement rates for the MMPI-2-RF F-r and Fp-r scales across ethnic, gender, and diagnostic groups with a forensic inpatient unit. *Psychological Assessment, 29*, 500-508. doi: [10.1037/pas0000366](https://doi.org/10.1037/pas0000366)
- Goodwin, B. E., Sellbom, M., & Arbisi, P. A. (2013). Post-traumatic stress disorder in veterans: The utility of the MMPI-2-RF Validity Scales in detecting over-reported symptoms. *Psychological Assessment, 25*, 671-678. doi: [10.1037/a0032214](https://doi.org/10.1037/a0032214)



- Greiffenstein, M., Gervais, R. O., Baker, W. J., Artiola, L., & Smith, H. (2013). Symptom validity testing in medically unexplained pain: A Chronic Regional Pain Syndrome Type 1 case series. *The Clinical Neuropsychologist*, 27, 138–147. doi: [10.1080/13854046.2012.722686](https://doi.org/10.1080/13854046.2012.722686)
- Grossi, L. M., Green, D., Einzig, S., & Belfi, B. (2017). Evaluation of the Response Bias Scale and the Improbable Failure Scale in assessing feigned cognitive impairment. *Psychological Assessment*, 29, 531–541. doi: [10.1037/pas0000364](https://doi.org/10.1037/pas0000364)
- Gu, W., Reddy, H.B., Green, D., Belfi, B., & Einzig, S. (2017). Inconsistent responding in a criminal forensic setting: An evaluation of the VRIN-r and TRIN-r scales of the MMPI-2-RF. *Journal of Personality Assessment*. doi: [10.1080/00223891.2016.1149483](https://doi.org/10.1080/00223891.2016.1149483)
- Handel, R. W., Ben-Porath, Y. S., Tellegen, A., & Archer, R. P. (2010). Psychometric functioning of the MMPI-2-RF VRIN-r and TRIN-r Scales with varying degrees of randomness, acquiescence, and counter-acquiescence. *Psychological Assessment*, 22, 87–95. doi: [10.1037/a0017061](https://doi.org/10.1037/a0017061)
- Haneveld, E. K., Kamphuis, J. H., Smid, W., & Forbey, J. D. (2016). Using MMPI-2-RF correlates to elucidate the PCL-R and its four facets in a sample of male forensic psychiatric patients. *Journal of Personality Assessment*. doi: [10.1080/00223891.2016.1228655](https://doi.org/10.1080/00223891.2016.1228655)
- Harp, J. P., Jasinski, L. J., Shandrea-Ochsner, A., Mason, L. H., & Berry, D. T. R. (2011). Detection of malingered ADHD using the MMPI-2-RF. *Psychological Injury and Law*, 4, 32–43. doi: [10.1007/s12207-011-9100-9](https://doi.org/10.1007/s12207-011-9100-9)
- Henry, G. K., Heilbronner, R. L., Algina, J., & Kaya, Y. (2013). Derivation of the MMPI-2-RF Henry-Heilbronner Index-r (HHI-r) Scale. *The Clinical Neuropsychologist*, 27, 509–515. doi: <http://dx.doi.org/10.1080/13854046.2012.739644>
- Henry, G. K., Heilbronner, R. L., Suhr, J., Gornbein, J., Wagner, E., & Drabe, D. L. (2018). Illness perceptions predict cognitive performance validity. *Journal of the International Neuropsychological Society*, 24, 1–11. doi: [10.1017/S1355617718000218](https://doi.org/10.1017/S1355617718000218)
- Hoeltzle, J. B., Nelson, N. W., & Arbisi, P. A. (2012). MMPI-2 and MMPI-2-Restructured Form Validity Scales: Complementary approaches to evaluate response validity. *Psychological Injury and Law*, 5, 174–191. doi: [10.1007/s12207-012-9139-2](https://doi.org/10.1007/s12207-012-9139-2)
- Ingram, P. B., & Ternes, M. S. (2016). The detection of content-based invalid responding: a meta-analysis of the MMPI-2-Restructured Form's (MMPI-2-RF) over-reporting validity scales. *The Clinical Neuropsychologist*, 30, 473–496 doi: [10.1080/13854046.2016.1187769](https://doi.org/10.1080/13854046.2016.1187769)
- Jiménez-Gómez, F., Sánchez-Crespo, G., & Ampudia-Rueda, A. (2013). Is there a social desirability scale in the MMPI-2-RF? *Clínica y Salud*, 24(3), 161–168. doi: [10.1016/S1130-5274\(13\)70017-3](https://doi.org/10.1016/S1130-5274(13)70017-3)
- Jones, A. (2016). Cutoff scores for MMPI-2 and MMPI-2-RF cognitive-somatic validity scales for psychometrically defined malingering groups in a military sample. *Archives of Clinical Neuropsychology*, 31, 76–81. doi: [10.1093/arclin/acw035](https://doi.org/10.1093/arclin/acw035)
- Jones, A., & Ingram, M. V. (2011). A comparison of selected MMPI-2 and MMPI-2-RF Validity Scales in assessing effort on cognitive tests in a military sample. *The Clinical Neuropsychologist*, 25, 1207–1227. doi: [10.1080/13854046.2011.600726](https://doi.org/10.1080/13854046.2011.600726)
- Jones, A., Ingram, M. V., & Ben-Porath, Y. S. (2012). Scores on the MMPI-2-RF Scales as a function of increasing levels of failure on cognitive symptom validity tests in a military sample. *The Clinical Neuropsychologist*, 26, 790–815. doi: [10.1080/13854046.2012.693202](https://doi.org/10.1080/13854046.2012.693202)

- Larrabee, G. J., Bianchini, K. J., Boone, K. B., Rohling, M. L. (2017). The MMPI-2/MMPI-2-RF Symptom Validity Scale (FBS/FBS-r) is not a measure of 'litigation response syndrome': commentary on Nichols and Gass (2015). *The Clinical Neuropsychologist*, 31, 1387-1389. doi: [10.1080/13854046.2017.1364423](https://doi.org/10.1080/13854046.2017.1364423)
- Larrabee, G. J., Bianchini, K. J., Boone, K. B., Rohling, M. L. (2017). The validity of the MMPI-2/MMPI-2-RF Symptom Validity Scale (FBS/FBS-r) is established: reply to Nichols (2017). *The Clinical Neuropsychologist*, 31, 1401-1405. doi: [10.1080/13854046.2017.1363293](https://doi.org/10.1080/13854046.2017.1363293)
- Lange, R. T., Brickell, T. A., & French, L. M. (2015). Examination of the Mild Brain Injury Atypical Symptom Scale and the Validity-10 Scale to detect exaggeration in US military service members. *Journal of Clinical and Experimental Neuropsychology*, 37, 325-337. doi: [10.1080/13803395.2015.1013021](https://doi.org/10.1080/13803395.2015.1013021)
- Lange, R. T., Sullivan, K. A., & Scott, C. (2010). Comparison of MMPI-2 and PAI validity indicators to detect feigned depression and PTSD symptom reporting. *Psychiatry Research*, 176, 229-235. doi: [10.1016/j.psychres.2009.03.004](https://doi.org/10.1016/j.psychres.2009.03.004)
- Lau, L., Basso, M. R., Estevis, E., Miller, A., Whiteside, D. M., Combs, D., & Arentsen, T. J. (2017). Detecting coached neuropsychological dysfunction: a simulation experiment regarding mild traumatic brain injury. *The Clinical Neuropsychologist*, 31, 1412-1431. doi: [10.1080/13854046.2017.1318954](https://doi.org/10.1080/13854046.2017.1318954)
- Manderino, L. M., & Gunstad, J. (2017). Performance of the Immediate Post-Concussion Assessment and Cognitive Testing protocol validity indices. *Archives of Clinical Neuropsychology*. doi: [10.1093/arclin/acx102](https://doi.org/10.1093/arclin/acx102)
- Marion, B. E., Sellbom, M., & Bagby, R. M. (2011). The detection of feigned psychiatric disorders using the MMPI-2-RF overreporting Validity Scales: An analog investigation. *Psychological Injury and Law*, 4, 1-12. doi: [10.1007/s12207-011-9097-0](https://doi.org/10.1007/s12207-011-9097-0)
- Marion, B. E., Sellbom, M., Salekin, R. T., Toomey, J. A., Kucharski, T., & Duncan, S. (2013). An examination of the association between psychopathy and dissimulation using the MMPI-2-RF Validity Scales. *Law and Human Behavior*, 37, 219-230. doi: [10.1037/lhb0000008](https://doi.org/10.1037/lhb0000008)
- Martin, P. K., Schroeder, R. W., Heinrichs, R. J., Baade, L. E. (2015). Does true neurocognitive dysfunction contribute to Minnesota Multiphasic Personality Inventory-2<sup>nd</sup> Edition-Restructured Form cognitive validity scores? *Archives of General Neuropsychology*. doi: [10.1093/arclin/acv032](https://doi.org/10.1093/arclin/acv032)
- Mason, L. H., Shandera-Ochsner, A. L., Williamson, K. D., Harp, J. P., Edmundson, M., Berry, D. T. R., & High, W. M. (2013). Accuracy of MMPI-2-RF Validity Scales for identifying feigned PTSD symptoms, random responding, and genuine PTSD. *Journal of Personality Assessment*, 95, 58-593. doi: [10.1080/00223891.2013.819512](https://doi.org/10.1080/00223891.2013.819512)
- McBride, W. F., Crighton, A. H., Wygant, D. B., & Granager, R. P. (2013). It's not all in your head (or at least your brain): Association of traumatic brain lesion presence and location with performance on measures of response bias in forensic evaluation. *Behavioral Sciences and the Law*, 31, 779-788. doi: [10.1002/bsl.2083](https://doi.org/10.1002/bsl.2083)
- McGee, S. A., Bagby, R. M., Goodwin, B. E., Burchett, D., Sellbom, M., Ayearst, L. E., Dhillon, S. Yiu, S., Ben-Porath, Y. S., & Baker, S. (2015). The effect of response bias on the Personality Inventory for DSM-5. *Journal of Personality Assessment*, 98, 51-61. doi: [10.1080/00223891.2015.1096791](https://doi.org/10.1080/00223891.2015.1096791)
- Meyers, J. E., Miller, R. M., Haws, N. A., Murphy-Tafti, J. L., Curtis, T. D., Rupp, Z. W., Smart, T. A., & Thompson, L. M. (2013). An adaptation of the MMPI-2 Meyers Index for the MMPI-2-RF. *Applied Neuropsychology: Adult*. doi: [10.1080/09084282.2013.780173](https://doi.org/10.1080/09084282.2013.780173)

- Nelson, N. W., Sweet, J. J., & Heilbronner, R. L. (2007). Examination of the new MMPI-2 Response Bias Scale (Gervais): Relationship with MMPI-2 Validity Scales. *Journal of Clinical and Experimental Neuropsychology*, 29, 67–72. doi: [10.1080/13803390500488546](https://doi.org/10.1080/13803390500488546)
- Nguyen, C. T., Green, D., & Barr, W. B. (2015). Evaluation of the MMPI-2-RF for detection of over-reported symptoms in civil forensic and disability setting. *The Clinical Neuropsychologist*, 29, 255–271. doi: [1080/13854046.2015.1033020](https://doi.org/1080/13854046.2015.1033020)
- Nichols, D. S. (2017). Fake Bad Scale: The case of the missing construct, a response to Larrabee, Bianchini, Boone, and Rohling (2017). *The Clinical Neuropsychologist*, 31, 1396-1400. doi: [10.1080/13854046.2017.1365934](https://doi.org/10.1080/13854046.2017.1365934)
- Nichols, D. S., & Gass, C. S. (2015). The Fake Bad Scale: Malingering or litigation response syndrome. Which is it? *Archives of Assessment Psychology*, 5, 5–10.
- Olsen, A.M., & Veltri, C. O. C. (2018). The moderating influence of disorder on coached overreporting using the MMPI-2-RF. *Journal of Personality Assessment*. doi: [10.1080/00223891.2018.1472099](https://doi.org/10.1080/00223891.2018.1472099)
- Pape, T. L., Harold, A. A., Smith, B., Babcock-Paziale, J., Harp, J., Shandera-Oschner, A., Jenjins, S., Evans, C. T., Schlembacker, R., & High, W. M. (2016). Algorithm for symptom attribution and classification following possible Mild Traumatic Injury. *Journal of Head Trauma Rehabilitation*. doi: [10.1097/HTR.0000000000000220](https://doi.org/10.1097/HTR.0000000000000220)
- Peck, C. P., Schroeder, R. W., Heinrichs, R. J., VonDran, E. J., Brockman, C. J., Webster, B. K., & Baade, L. E. (2013). Differences in MMPI-2 FBS and RBS scores in brain injury, probable malingering, and conversion disorder groups: A preliminary study. *The Clinical Neuropsychologist*, 27, 693–707. doi: [10.1080/13854046.2013.779032](https://doi.org/10.1080/13854046.2013.779032)
- Ray, C. L. (2017). Practical use of the MMPI-2-RF validity indicators in VA compensation and pension evaluations. *Psychological Injury and Law*. doi: [10.1007/s12207-017-9289-3](https://doi.org/10.1007/s12207-017-9289-3)
- Robinson, E. V., & Rogers, R. (2017). Detection of feigned ADHD across two domains: MMPI-2-RF and CAARS for faked symptoms and TOVA for simulated attention deficits. *Journal of Psychopathology and Behavioral Assessment*. doi: [10.1007/s10862-017-9640-8](https://doi.org/10.1007/s10862-017-9640-8)
- Rogers, R., Gillard, N., Berry, D. T. R., & Granacher, R. P. (2011). Effectiveness of the MMPI-2-RF Validity Scales for feigned mental disorders and cognitive impairment: A known-groups study. *Journal of Psychopathology and Behavioral Assessment*, 33, 355–367. doi: [10.1007/s10862-011-9222-0](https://doi.org/10.1007/s10862-011-9222-0)
- Roma, P., Verrocchio, M. C., Mazza, C., Marchetti, D., Burla, F., Cinti, M. E., & Ferracuti, S. (2018). Could time detect a faking-good attitude? A study with the MMPI-2-RF. *Frontiers in Psychology*. doi: [10.3389/fpsyg.2018.01064](https://doi.org/10.3389/fpsyg.2018.01064)
- Sanchez, G., Ampudia, A. Jimenez, F., & Amado, B. G. (2017). Contrasting the efficacy of the MMPI-2-RF overreporting scales in the detection of malingering. *The European Journal of Psychology Applied in the Legal Context*, 9, 51-56. doi: [10.1016/j.ejpal.2017.03.002](https://doi.org/10.1016/j.ejpal.2017.03.002)
- Sharf, A. J., Rogers, R., Williams, M. M., & Henry, S. A. (2017). The effectiveness of the MMPI-2-RF in detecting feigned mental disorders and cognitive deficits: A meta-analysis. *Journal of Psychopathology and Behavioral Assessment*, 39, 441-455. doi: [10.1007/s10862-017-9590-1](https://doi.org/10.1007/s10862-017-9590-1)
- Schroeder, R. W., Baade, L. E., Peck, C. P., VonDran, E., Brockman, C. J., Webster, B. K., & Heinrichs, R. J. (2012). Validation of the MMPI-2-RF Validity Scales in criterion group neuropsychological samples. *The Clinical Neuropsychologist*, 26, 129–146. doi: [10.1080/13854046.2011.639314](https://doi.org/10.1080/13854046.2011.639314)

- Sellbom, M., & Bagby, R. M. (2008). Validity of the MMPI-2-RF (Restructured Form) L-r and K-r Scales in detecting underreporting in clinical and nonclinical samples. *Psychological Assessment, 20*, 370–376. doi: [10.1037/a0012952](https://doi.org/10.1037/a0012952)
- Sellbom, M., & Bagby, R. M. (2010). The detection of over-reported psychopathology with the MMPI-2-RF (Restructured Form) Validity Scales. *Psychological Assessment, 22*, 757–767. doi: [10.1037/a0020825](https://doi.org/10.1037/a0020825)
- Sellbom, M., Dhillon, S., & Bagby, R. M. (2018). Development and validation of an overreporting scale for the Personality Inventory for DSM-5 (PID-5). *Psychological Assessment, 30*, 582-593. doi: [10.1037/pas0000507](https://doi.org/10.1037/pas0000507)
- Sellbom, M., Toomey, A., Wygant, D., Kucharski, L. T., & Duncan, S. (2010). Utility of the MMPI-2-RF (Restructured Form) Validity Scales in detecting malingering in a criminal forensic setting: A known-groups design. *Psychological Assessment, 22*, 22–31. doi: [10.1037/a0018222](https://doi.org/10.1037/a0018222)
- Sellbom, M., Wygant, D. B., & Bagby, R. M. (2012). Utility of the MMPI-2-RF in detecting non-credible somatic complaints. *Psychiatry Research, 197*, 295–301. doi: [10.1016/j.psychres.2011.12.043](https://doi.org/10.1016/j.psychres.2011.12.043)
- Shkalim, E., Ben-Porath, Y. S., Handel, R. W., Almagor, M., & Tellegen, A. (2016). Psychometric examination, adaptation, and evaluation of the Hebrew translation of the MMPI-2-RF VRIN-r and TRIN-r Validity Scales. *Journal of Personality Assessment, 98*, 608-615. doi: [10.1080/00223891.2016.1174705](https://doi.org/10.1080/00223891.2016.1174705)
- Shura, R. D., Denning, J. H., Miskey, H. M., & Rowland, J. A. (2017). Symptom and performance validity with veterans assessed for Attention-Deficit/Hyperactivity Disorder. *Psychological Assessment, 29*, 1458-1465. doi: [10.1037/pas0000436](https://doi.org/10.1037/pas0000436)
- Sleep, C., Sellbom, M., Campbell, W. K., & Miller, J. D. (2016). Narcissism and response validity: Do narcissistic individuals under report psychopathology? *Psychological Assessment*. doi: [10.1037/pas0000413](https://doi.org/10.1037/pas0000413)
- Smart, C. M., Nelson, N. W., Sweet, J. J., Bryant, F. B., Berry, D. T. R., Granacher, R. P., & Heilbronner, R. L. (2008). Use of MMPI-2 to predict cognitive effort: A hierarchically optimal classification tree analysis. *Journal of the International Neuropsychological Society, 14*, 842–852. doi: [10.1017/S1355617708081034](https://doi.org/10.1017/S1355617708081034)
- Sullivan, K. A. & Elliott, C. (2012). An investigation of the validity of the MMPI-2 Response Bias Scale using an analog simulation design. *The Clinical Neuropsychologist, 26*, 160–176. doi: [10.1080/13854046.2011.647084](https://doi.org/10.1080/13854046.2011.647084)
- Sullivan, K. A., Elliott, C. D., Lange, R. T., & Anderson, D. S. (2013). A known groups evaluation of the Response Bias Scale in a neuropsychological setting. *Applied Neuropsychology: Adult, 20*, 20–32. doi: [10.1080/09084282.2012.670149](https://doi.org/10.1080/09084282.2012.670149)
- Tarescavage, A., Wygant, D. B., Gervais, R. O., & Ben-Porath, Y. S. (2013). Association between the MMPI-2 Restructured Form (MMPI-2-RF) and malingered neurocognitive dysfunction among non-head injury disability claimants. *The Clinical Neuropsychologist, 27*, 313–335. doi: [10.1080/13854046.2012.744099](https://doi.org/10.1080/13854046.2012.744099)
- Tolin, D. F., Steenkamp, M. M., Marx, B. P., & Litz, B. T. (2010). Detecting symptom exaggeration in combat veterans using the MMPI-2 symptom Validity Scales: A mixed group validation. *Psychological Assessment, 22*, 729–736. doi: [10.1037/a0020973](https://doi.org/10.1037/a0020973)
- Tsushima, W. T., Geling, O., & Fabriga, L. (2011). Comparison of MMPI-2 Validity Scale scores of personal injury litigants and disability claimants. *The Clinical Neuropsychologist, 25*, 1403–1414. doi: [10.1080/13854046.2011.613854](https://doi.org/10.1080/13854046.2011.613854)



- Tylicki, J. L., Wygant, D. B., Tarescavage, A. M., Frederick, R. J., Tyner, E. A., Grannacher, R. P., & Sellbom, M. (2018). Comparability of Structured Interview of Reported Symptoms (SIRS) and Structured Interview of Reported Symptom-2nd Edition (SIRS-2) classifications with external response bias criteria. *Psychological Assessment*. doi: [10.1037/pas0000573](https://doi.org/10.1037/pas0000573)
- Van Dyke, S. A., Axelrod, B. N., & Schutte, C. (2010). The utility of the Post-concussive Syndrome Questionnaire. *Archives of Clinical Neuropsychology*, 25, 634–639. doi: [10.1093/arclin/acq063](https://doi.org/10.1093/arclin/acq063)
- Van Dyke, S. A., Millis, S. R., Axelrod, B. N., & Hanks, R. A. (2013). Assessing effort: Differentiating performance and symptom validity. *The Clinical Neuropsychologist*. doi: [10.1080/13854046.2013.835447](https://doi.org/10.1080/13854046.2013.835447)
- Wall, T. D., Wygant, D. B., & Gallagher, R. W. (2015). Identifying over-reporting in a correctional setting: Utility of the MMPI-2 Restructured Form Validity Scales. *Criminal Justice and Behavior*, 42, 610–622. doi: [10.1177/0093854814556881](https://doi.org/10.1177/0093854814556881)
- Wershba, R. E., Locke, D. E. C., & Lanyon, R. I. (2015). Analysis of Minnesota Multiphasic Personality Inventory-2-Restructured Form bias indicators as suppressor or moderators in a medical setting. *Psychological Assessment*, 27, 733–737. doi: [10.1037/a0038802](https://doi.org/10.1037/a0038802)
- Whitney, K. W. (2013). Predicting test of memory malingering and medical symptom validity test failure within a veterans affairs medical center: Use of the Response Bias Scale and the Henry-Heilbronner Index. *Archives of Clinical Neuropsychology*, 28, 222–235. doi: [10.1093/arclin/act012](https://doi.org/10.1093/arclin/act012)
- Whitney, K. A., Davis, J. J., Shephard, P. H., & Herman, S. M. (2008). Utility of the Response Bias Scale and other MMPI-2 Validity Scales in predicting TOMM performance. *Archives of Clinical Neuropsychology*, 23, 777–786. doi: [10.1016/j.acn.2008.09.001](https://doi.org/10.1016/j.acn.2008.09.001)
- Wiggins, C. W., Wygant, D. B., Hoelzle, J. B., & Gervais, R. O. (2012). The more you say the less it means: Over-reporting and attenuated criterion validity in a forensic disability sample. *Psychological Injury and Law*, 5, 162–173. doi: [10.1007/s12207-012-9137-4](https://doi.org/10.1007/s12207-012-9137-4)
- Wygant, D. B., Anderson, J. L., Sellbom, M., Rapier, J. L., Algeier, L. M., & Granacher, R. P. (2011). Association of MMPI-2 Restructured Form (MMPI-2-RF) Validity Scales with structured malingering criteria. *Psychological Injury and Law*, 4, 13–23. doi: [10.1007/s12207-011-9098-z](https://doi.org/10.1007/s12207-011-9098-z)
- Wygant, D. B., Arbisi, P. A., Bianchini, B. J., Umlauf, R.L. (2016). Waddell nonorganic signs: New evidence suggests somatic amplification among outpatient chronic pain patients. *The Spine Journal*. doi: [10.1016/j.spinee.2016.10.018](https://doi.org/10.1016/j.spinee.2016.10.018)
- Wygant, D. B., Ben-Porath, Y. S., Arbisi, P. A., Berry, D. T. R., Freeman, D. B., & Heilbronner, R. L. (2009). Examination of the MMPI-2 Restructured Form (MMPI-2-RF) Validity Scales in civil forensic settings: Findings from simulation and known group samples. *Archives of Clinical Neuropsychology*, 27, 671–680. doi: [10.1093/arclin/acp073](https://doi.org/10.1093/arclin/acp073)
- Wygant, D. B., Sellbom, M., Gervais, R. O., Ben-Porath, Y. S., Stafford, K. P., Freeman, D. B., & Heilbronner, R. L. (2010). Further validation of the MMPI-2 and MMPI-2-RF Response Bias Scale: Findings from disability and criminal forensic settings. *Psychological Assessment*, 22, 745–756. doi: [10.1037/a0020042](https://doi.org/10.1037/a0020042)
- Young, J. C., & Gross, A. M. (2011). Detection of response bias and noncredible performance in adult Attention-Deficit/Hyperactivity Disorder. *Archives of Clinical Neuropsychology*, 26, 165–175. doi: [10.1093/arclin/acr013](https://doi.org/10.1093/arclin/acr013)



Young, J. C., Kearns, L. A., & Roper, B. L. (2011). Validation of the MMPI-2 Response Bias Scale and Henry-Heilbronner Index in a U.S. veteran population. *Archives of Clinical Neuropsychology*, 3, 194–204. doi: [10.1093/arclin/acr015](https://doi.org/10.1093/arclin/acr015)

Youngjohn, J. R., Weshba, R., Stevenson, M., Sturgeon, J., & Thomas, M. L. (2011). Independent validation of the MMPI-2-RF Somatic/Cognitive and Validity Scales in TBI litigants tested for effort. *The Clinical Neuropsychologist*, 25, 463–476. doi: [10.1080/13854046.2011.554444](https://doi.org/10.1080/13854046.2011.554444)

## CORRECTIONAL SETTINGS

Clegg, C., Fremouw, W., Horacek, T., Cole, A., & Schwartz, R. (2010). Factors associated with treatment acceptance and compliance among incarcerated male sex offenders. *International Journal of Offender Therapy and Comparative Criminology*, 6, 880–897. doi: [10.1177/0306624X10376160](https://doi.org/10.1177/0306624X10376160)

Forbey, J. D., Ben-Porath, Y. S., & Gartland, D. (2009). Validation of the MMPI-2 Computerized Adaptive Version (MMPI-2-CA) in a correctional intake facility. *Psychological Services*, 6, 279–292. doi: [10.1037/a0016195](https://doi.org/10.1037/a0016195)

Glenn, A. R., Sellbom, M. (2015). Theoretical and empirical concerns regarding the dark triad as a construct. *Journal of Personality Disorders*, 29:3, 360–377. doi: [10.1521/pedi\\_2014\\_28\\_162](https://doi.org/10.1521/pedi_2014_28_162)

Gotfried, E. D., Anestis, J. C., Dillon, K. H., & Carbonell, J. J. (2016). The associations between Minnesota Multiphasic Personality Inventory-2-Restructured Form and self-reported physical and sexual abuse and posttraumatic symptoms in a sample of incarcerated women. *International Journal of Forensic Mental Health*. doi: [10.1080/14999013.2016.1228088](https://doi.org/10.1080/14999013.2016.1228088)

Johnson, A. K., Sellbom, M., & Phillips, T. R. (2014). Elucidating the associations between psychopathy, Gray's BIS and BAS constructs, and externalizing behavior. *Personality and Individual Differences*. doi: [10.1016/j.paid.2014.06.026](https://doi.org/10.1016/j.paid.2014.06.026)

Kastner, R. M., Sellbom, M., & Lilienfeld, S. O. (2012). A comparison of the psychometric properties of the Psychopathic Personality Inventory Full-Length and Short-Form Versions. *Psychological Assessment*, 24, 261–267. doi: [10.1037/a0025832](https://doi.org/10.1037/a0025832)

Kutchen, T. J., Wygant, D. B., Tylicki, J. L., Dieter, A. M., Veltri, C. O., & Sellbom, M. (2017). Construct validity of the MMPI-2-RF Triarchic Psychopathy Scales in correctional and collegiate samples. *Journal of Personality Assessment*, 99, 408–415. doi: [10.1080/00223891.2016.1238829](https://doi.org/10.1080/00223891.2016.1238829)

Laurinaitytė, I., Laurinavičius, A., Ustinavičiūtė, L., Wygant, D. B., Sellbom, M. (2017). Utility of the MMPI-2 Restructured Form (MMPI-2-RF) in a sample of Lithuanian Male Offenders. *Law and Human Behavior*, 41, 494–505. doi: [10.1037/lhb0000254](https://doi.org/10.1037/lhb0000254)

McAnulty, R. D., McAnulty, D. P., Sipp, J. E., Demakis, G. J., & Heggstad, E. D. (2014). Predictive validity of the MMPI-2 among female offenders in a residential treatment program. *Journal of Personality Assessment*, 99, 341–350. doi: [10.1080/00223891.2014.880061](https://doi.org/10.1080/00223891.2014.880061)

Phillips, T. R., Sellbom, M., Ben-Porath, Y. S., & Patrick, C. J. (2013). Further development and construct validation of MMPI-2-RF indices of global psychopathy, fearless-dominance, and impulsive-antisociality in a sample of incarcerated women. *Law and Human Behavior*, 38, 34–46. doi: [10.1037/lhb0000040](https://doi.org/10.1037/lhb0000040)

Sellbom, M. (2017). Mapping the MMPI-2-RF Specific Problems scales onto Extant Psychopathology Structures. *Journal of Personality Assessment*. doi: [10.1080/00223891.2016.1206909](https://doi.org/10.1080/00223891.2016.1206909)

- Sellbom, M. (2014). A factor mixture model approach to elaborating on offender mental health classification with the MMPI-2-RF. *Journal of Personality Assessment, 96*, 293–305. doi: [10.1080/00223891.2013.843538](https://doi.org/10.1080/00223891.2013.843538)
- Sellbom, M. (2011). Elaborating on the construct validity of the Levenson Self-Report Psychopathy Scale in incarcerated and non-incarcerated samples. *Law and Human Behavior, 35*, 440–451. doi: [10.1007/s10979-010-9249-x](https://doi.org/10.1007/s10979-010-9249-x)
- Sellbom, M., Ben-Porath, Y. S., Patrick, C. J., Wygant, D. B., Gartland, D. M., & Stafford, K. P. (2012). Development and construct validation of MMPI-2-RF measures assessing global psychopathy, fearless-dominance, and impulsive-antisociality. *Personality Disorders: Theory, Research, and Treatment, 3*, 17–38. doi: [10.1037/a0023888](https://doi.org/10.1037/a0023888)
- Sellbom, M., Donnelly, K., Rock, R. C., Phillips, T. R., & Ben-Porath, Y. S. (2017). Examining gender as moderating the association between psychopathy and substance abuse. *Psychology, Crime, and Law, 23*, 376–390. doi: [10.1080/1068316X.2016.1258466](https://doi.org/10.1080/1068316X.2016.1258466)
- Sellbom, M., Drislane, L. E., Johnson, A. K., Goodwin, B. E., Phillips, T. R., & Patrick, C. J. (2017). Development and validation of MMPI-2-RF scales for indexing triarchic psychopathy constructs. *Assessment, 23*, 527–543. doi: [10.1177/1073191115590853](https://doi.org/10.1177/1073191115590853)

### FORENSIC SETTINGS

- Acre, R., Farina, F., Seijo, D., & Novo, M. (2015). Assessing Impression Management with the MMPI-2 in child custody litigation. *Assessment, 22*, 769–777. doi: [10.1177/1073191114558111](https://doi.org/10.1177/1073191114558111)
- Archer, E. M., Hagan, L. D., Mason, J., Handel, R. W., & Archer, R. P. (2012). MMPI-2-RF characteristics of custody evaluation litigants. *Assessment, 19*, 14–20. doi: [10.1177/1073191110397469](https://doi.org/10.1177/1073191110397469)
- Anderson, J. L., Sellbom, M., Pymont, C., Smid, W., De Saeger, H. & Kamphuis, J. H. (2015). Measurement of DSM-5 Section II personality disorder constructs using the MMPI-2-RF in clinical and forensic samples. *Psychological Assessment, 27*, 786–800. doi: [10.1037/pas0000103](https://doi.org/10.1037/pas0000103)
- Anderson, J. L., Wood, M. E., Tarescavage, A. M., Burchette, D., & Glassmire, D. M. (2017). The role of dimensional personality psychopathology in a forensic inpatient psychiatric setting. *Journal of Personality Disorders*. doi: [10.1521/pedi\\_2017\\_31\\_301](https://doi.org/10.1521/pedi_2017_31_301)
- Balasanya, M., Boone, K. B., Ermshar, A., Miora, D., Cottingham, M., Ziegler, E., Zeller, M. A., Wright, M. (2017). Examination of the Modified Somatic Perception Questionnaire (MSPQ) in a large sample of credible and noncredible patients referred for neuropsychological testing. *The Clinical Neuropsychologist, 32*, 165–182. doi: [10.1080/13854046.2017.1330421](https://doi.org/10.1080/13854046.2017.1330421)
- Ben-Porath, Y. S. (2012). Addressing challenges to MMPI-2-RF-based testimony: Questions and answers. *Archives of Clinical Neuropsychology, 27*, 691–705. doi: [10.1093/arclin/acs083](https://doi.org/10.1093/arclin/acs083)
- Ben-Porath, Y. S. (2018). Uses and misuses of Ted Kaczynski's MMPI. *Journal of Personality Assessment*. doi: [10.1080/00223891.2018.1468337](https://doi.org/10.1080/00223891.2018.1468337)
- Ben-Porath, Y. S. & Flens, J. R. (2012). Butcher and Williams's (this issue) critique of the MMPI-2-RF is slanted and misleading. *Journal of Child Custody, 9*, 223–232. doi: [10.1080/15379418.2012.748605](https://doi.org/10.1080/15379418.2012.748605)
- Butcher, J. N., & Williams, C. L. (2012). Problems with using the MMPI-2-RF in forensic evaluations: A clarification to Ellis. *Journal of Child Custody, 9*, 217–222. doi: [10.1080/15379418.2012.748347](https://doi.org/10.1080/15379418.2012.748347)

- Downing, S. K., Denney, R. L., Spray, B. J., Houston, C. M., & Halfaker, D. H. (2008). Examining the relationship between the Restructured Scales and the Fake Bad Scale of the MMPI-2. *The Clinical Neuropsychologist, 22*, 680–688. doi: [10.1080/13854040701562825](https://doi.org/10.1080/13854040701562825)
- Gervais, R. O., Ben-Porath, Y. S., & Wygant, D. B. (2009). Empirical correlates and interpretation of the MMPI-2-RF Cognitive Complaints Scale. *The Clinical Neuropsychologist, 23*, 996–1015. doi: [10.1080/13854040902748249](https://doi.org/10.1080/13854040902748249)
- Glassmire, D. M., Tarescavage, A. M., Burchett, D., Martinez, J., & Gomez, A. (2016). Clinical utility of the MMPI-2-RF SUI items and scale in a forensic inpatient setting: Association with interview self-reports and future suicidal behavior. *Psychological Assessment, 28*, 1502-1509. doi: [doi.org/10.1037/pas0000220](https://doi.org/10.1037/pas0000220)
- Grossi, L. M., Green, D., Belfi, B., McGrath, R.E., Griswald, H., & Schreiber, J. (2015). Identifying aggression in forensic inpatients using the MMPI-2-RF: An examination of MMPI-2-RF scale scores and estimated psychopathy indices. *International Journal of Forensic Mental Health, 14*, 231–244. doi: [10.1080/14999013.2015.1108943](https://doi.org/10.1080/14999013.2015.1108943)
- Grossi, L. M., Green, D., Schneider, M., Belfi, B., & Segal, S. (2018). Personality, psychiatric, and cognitive predictors of length of time for competency to stand trial restoration. *International Journal of Forensic Mental Health, 17*, 167-180. doi: [10.1080/14999013.2018.1459964](https://doi.org/10.1080/14999013.2018.1459964)
- Henry, G. K., Heilbronner, R. L., Mittenberg, W., Enders, C., & Dombaski, K. (2009). Comparison of the MMPI-2 Restructured Demoralization Scale, Depression Scale, and Malingered Mood Disorder Scale in identifying non-credible symptom reporting in personal injury litigants and disability claimants. *The Clinical Neuropsychologist, 23*, 153–166. doi: [10.1080/13854040801969524](https://doi.org/10.1080/13854040801969524)
- Henry, G. K., Heilbronner, R. L., Mittenberg, W., Enders, C., & Stanzal, S. R. (2008). Comparison of the Lees-Haley Fake Bad Scale, Henry-Heilbronner Index, and Restructured Clinical Scale 1 in identifying noncredible symptom reporting. *The Clinical Neuropsychologist, 22*, 919–929. doi: [10.1080/13854040701625853](https://doi.org/10.1080/13854040701625853)
- Himsl, K., Burchett, D., Tarescavage, A. M., & Glassmire, D. M. (2017). Assessing reading ability for psychological testing in forensic assessments: An investigation with the WRAT-4 and MMPI-2-RF. *International Journal of Forensic Mental Health, 16*, 239-248. doi: [10.1080/14999013.2017.1330293](https://doi.org/10.1080/14999013.2017.1330293)
- Kauffman, C. M., Stolberg, R., & Madero, J. (2015). An examination of the MMPI-2-RF (Restructured Form) with the MMPI-2 and MCMI-III in child custody litigants. *Journal of Child Custody, 12*, 129–151. doi: [10.1080/15379418.2015.1057354](https://doi.org/10.1080/15379418.2015.1057354)
- Klein Haneveld, E., Kamphuis, J. H., Smid, W., & Forbey, J. D. (2017). Using MMPI-2-RF correlates to elucidate the PCL-R and its four facets in a sample of male forensic psychiatric patients. *Journal of Personality Assessment, 99*, 398-407. doi: [10.1080/00223891.2016.1228655](https://doi.org/10.1080/00223891.2016.1228655)
- Lawson, A. K., Wright, C. V., & Fitzgerald, L. F. (2013). The evaluation of sexual harassment litigants: Reducing discrepancies in the diagnosis of post-traumatic stress disorder. *Law and Human Behavior, 37*, 337–347. doi: [10.1037/lhb0000024](https://doi.org/10.1037/lhb0000024)
- Mattson, C. A., Powers, B. K., Halfaker, D., Akenson, S. T., & Ben-Porath, Y. S. (2012). Predicting drug court completion with the MMPI-2-RF. *Psychological Assessment, 24*, 937–943. doi: [10.1037/a0028267](https://doi.org/10.1037/a0028267)
- Pinsooneault, T. B., & Ezzo, F. R. (2012). A comparison of MMPI-2-RF profiles between child maltreatment and non-maltreatment custody cases. *Journal of Forensic Psychology Practice, 12*, 227–237. doi: [10.1080/15228932.2012.674469](https://doi.org/10.1080/15228932.2012.674469)

- Resendes, J., & Lecci, L. (2012). Comparing the MMPI-2 scale scores of parents involved in parental competency and child custody assessments. *Psychological Assessment, 24*, 1054–1059. doi: [10.1037/a0028585](https://doi.org/10.1037/a0028585)
- Rock, R. C., Sellbom, M., Ben-Porath, Y. S., & Salekin, R. T. (2013). Concurrent and predictive validity of psychopathy in a batterers intervention sample. *Law and Human Behavior, 37*, 145–154. doi: [10.1037/lhb0000006](https://doi.org/10.1037/lhb0000006)
- Romero, I. E., Toorabally, N., Burchett, D., Tarescavage, A. M., & Glassmire, D. M. (2017). Mapping the MMPI-2-RF substantive scales onto, internalizing, externalizing, and thought dysfunction dimensions in a forensic inpatient setting. *Journal of Personality Assessment, 99*, 351–362. doi: [10.1080/00223891.2016.1223681](https://doi.org/10.1080/00223891.2016.1223681)
- Sellbom, M. (2012). The MMPI-2-RF is ready for the Daubert challenge: Evidence, implications, and recommendations for use in court testimony. *Journal of Psychological Practice, 17*, 151–179.
- Sellbom, M. (2016). Elucidating the validity of the externalizing spectrum of psychopathology in correctional, forensic, and community Samples. *Journal of Abnormal Psychology, 125*, 1027–1038. doi: [10.1037/abn0000171](https://doi.org/10.1037/abn0000171)
- Sellbom, M. (2017). Using the MMPI-2-RF to Characterize Defendants Evaluated for Competency to Stand Trial and Criminal Responsibility. *International Journal of Forensic Mental Health, 16*, 304–312. doi: [10.1080/14999013.2017.1371259](https://doi.org/10.1080/14999013.2017.1371259)
- Sellbom, M., & Bagby, R. M. (2009). Identifying PTSD personality subtypes in a workplace trauma sample. *Journal of Traumatic Stress, 22*, 471–475. doi: [10.1002/jts.20452](https://doi.org/10.1002/jts.20452)
- Sellbom, M., Ben-Porath, Y. S., Baum, L. J., Erez, E., & Gregory, C. (2008). Predictive validity of the MMPI-2 Restructured Clinical (RC) Scales in a batterers' intervention program. *Journal of Personality Assessment, 90*, 129–135. doi: [10.1080/00223890701845153](https://doi.org/10.1080/00223890701845153)
- Sellbom, M., & Ben-Porath, Y. S., & Stafford, K. P. (2007). A comparison of measures of psychopathic deviance in a forensic setting. *Psychological Assessment, 19*, 430–436. doi: [10.1037/1040-3590.19.4.430](https://doi.org/10.1037/1040-3590.19.4.430)
- Sellbom, M., Lee, T. T. C., Ben-Porath, Y. S., Arbisi, P. A., & Gervais, R. O. (2012). Differentiating PTSD symptomatology with the MMPI-2-RF (Restructured Form) in a forensic disability sample. *Psychiatry Research, 197*, 172–179. doi: [10.1016/j.psychres.2012.02.003](https://doi.org/10.1016/j.psychres.2012.02.003)
- Sellbom, M., Smid, W., De Saeger, H., Smit, N., & Kamphuis, J. H. (2014). Mapping the Personality Psychopathology Five domains onto DSM-IV personality disorders in Dutch clinical and forensic samples: Implications for the DSM-5. *Journal of Personality Assessment, 96*, 185–191. doi: [10.1080/00223891.2013.825625](https://doi.org/10.1080/00223891.2013.825625)
- Sleep, C. C., Petty, J. A., & Wygant, D. B. (2015). Framing the results: Assessment of response bias through select self-report measures in psychological injury evaluations. *Psychological Injury and Law, 8*, 27–309. doi: [10.1007/s12207-015-9219-1](https://doi.org/10.1007/s12207-015-9219-1)
- Solomon, D., Morgan, B., Asberg, K., & Achee, D. (2014). Treatment implications based on measures of child abuse potential and parental mental health: Are we missing an intervention opportunity? *Children and Youth Services Review, 43*, 153–159. doi: [10.1016/j.childyouth.2014.05.016](https://doi.org/10.1016/j.childyouth.2014.05.016)
- Stredny, R. V., Archer, R. P., & Mason, J. A. (2006). MMPI-2 and MCMI-III characteristics of parental competency examinees. *Journal of Personality Assessment, 87*, 113–115. doi: [10.1207/s15327752jpa8701\\_10](https://doi.org/10.1207/s15327752jpa8701_10)

- Tarescavage, A. M., Alosco, M. L., Ben-Porath, Y. S., Wood, A., & Luna-Jones, L. (2015). MMPI-2-RF scores generated from the MMPI-2 and MMPI-2-RF test booklets: Comparability in a sample of criminal defendants. *Assessment, 22*, 188-197. doi: [10.1177/1073191114537347](https://doi.org/10.1177/1073191114537347)
- Tarescavage, A. M., Cappo, B. M., & Ben-Porath, Y. S. (2018). Assessment of sex offenders with the Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF). *Sexual Abuse: A Journal of Research and Treatment, 30*, 413-437. doi: [10.1177/1079063216667921](https://doi.org/10.1177/1079063216667921)
- Tarescavage, A. M., Glassmire, D. M., & Burchett, D. (2016). Introduction of a conceptual model for integrating the MMPI-2-RF into HCR-20V3 violence risk assessments and associations between the MMPI-2-RF and institutional violence. *Law and Human Behavior, 40*, 626-637. doi: [10.1037/lhb0000207](https://doi.org/10.1037/lhb0000207)
- Tarescavage, A. M., Glassmire, D. M., & Burchett, D. (2018). Minnesota Multiphasic Personality Inventory-2-Restructured Form markers of future suicidal behavior in a forensic psychiatric hospital. *Psychological Assessment, 30*, 170-178. doi: [10.1037/pas0000463](https://doi.org/10.1037/pas0000463)
- Tarescavage, A. M., Luna-Jones, L., & Ben-Porath, Y. S. (2014). Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) predictors of violating probation after felonious crimes. *Psychological Assessment, 26*, 1375-1380. doi: [10.1037/pas0000022](https://doi.org/10.1037/pas0000022)
- Thomas, M. L., & Youngjohn, J. R. (2009). Let's not get hysterical: Comparing the MMPI-2 Validity, Clinical, and RC Scales in TBI litigants tested for effort. *The Clinical Neuropsychologist, 23*, 1067-1084. doi: [10.1080/13854040902795000](https://doi.org/10.1080/13854040902795000)
- Vines, L. M., Wygant, D. B., & Gervais, R. O. (2012). Empirically guided case conceptualization of posttraumatic stress disorder with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) in a forensic disability evaluation. *Journal of Psychological Practice, 17*, 180-205.
- Wise, E. A. (2009). Selected MMPI-2 scores of forensic offenders in a community setting. *Journal of Forensic Psychology Practice, 9*, 299-309. doi: [10.1080/15228930902936048](https://doi.org/10.1080/15228930902936048)
- Wolf, E. J., & Miller, M. W. (2014). The Minnesota Multiphasic Personality Inventory-2 Restructured Form and posttraumatic stress disorder: Forensic applications and considerations. *Psychological Injury and Law, 7*, 143-152. doi: [10.1007/s12207-014-9193-z](https://doi.org/10.1007/s12207-014-9193-z)
- Wygant, D. B., & Granacher, R. P. (2015). Assessment of validity and response bias in neuropsychiatric evaluations. *NeuroRehabilitation, 36*, 427-438.
- Wygant, D. B., & Sellbom, M. (2012). Viewing psychopathy from the perspective of the Personality Psychopathology Five model: Implications for DSM-5. *Journal of Personality Disorders, 26*, 717-726. doi: [10.1521/pedi.2012.26.5.717](https://doi.org/10.1521/pedi.2012.26.5.717)
- Wygant, D. B., Sellbom, M., Ben-Porath, Y. S., Stafford, K. P., Freeman, D. B., & Heilbronner, R. I. (2007). The relation between symptom validity testing and MMPI-2 scores as a function of forensic evaluation context. *Archives of Clinical Neuropsychology, 22*, 488-499. doi: [10.1016/j.acn.2007.01.027](https://doi.org/10.1016/j.acn.2007.01.027)

## MEDICAL SETTINGS

- Aguerrevere, L. E., Calamia, M. R., Greve, K. W., Bianchini, K. J., & Curtis, K. L. (2018). Clusters of financially incentivized chronic pain patients using the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF). *Psychological Assessment, 30*, 634-644. doi: [10.1037/pas0000509](https://doi.org/10.1037/pas0000509)



- Block, A. R., Ben-Porath, Y. S., & Marek, R. J. (2018). Patient activation mediates associations between psychosocial risk factors and spine surgery results. *Journal of Clinical Psychology in Medical Settings*. doi: [10.1007/s10880-018-9571-x](https://doi.org/10.1007/s10880-018-9571-x)
- Block, A. R., Ben-Porath, Y. S., & Marek, R. J. (2013). Psychological risk factors for poor outcome of spine surgery and spinal cord stimulator implant: A review of the literature and their assessment with the MMPI-2-RF. *The Clinical Neuropsychologist*, 27, 81–107 doi: [10.1080/13854046.2012.721007](https://doi.org/10.1080/13854046.2012.721007)
- Block, A. R., Marek, R. J., Ben-Porath, Y. S., & Kukal, D. (2017). Associations between pre-implant psychosocial factors and spinal cord stimulation outcome: Evaluation using the MMPI-2-RF. *Assessment*, 24, 60–70. doi: [10.1177/1073191115601518](https://doi.org/10.1177/1073191115601518)
- Block, A. R., Marek, R. J., Ben-Porath, Y. S., & Ohnmeiss, D. D. (2014). Associations between MMPI-2-RF scores, workers' compensation status, and spine surgery outcome. *Journal of Applied Biobehavioral Research*, 19, 248–267. doi: [10.1111/jabr.12028](https://doi.org/10.1111/jabr.12028)
- Block, A., Ohnmeiss, D., Ben-Porath, Y. S., & Burchett, D. (2011). Presurgical psychological screening: A new algorithm, including the MMPI-2-RF, for predicting surgery results. *Proceedings of the NASS 26th Annual Meeting/The Spine Journal* 11 (2011) 15–173S. doi: [10.1016/j.spinee.2011.08.333](https://doi.org/10.1016/j.spinee.2011.08.333)
- Bowden, S. C., White, J. R., Simpson, L., & Ben-Porath, Y. S. (2014). Elevation discrepancies between MMPI-2 Clinical and MMPI-2-RF Restructured Clinical (RC) Scales in people with seizure disorders. *Epilepsy and Behavior*, 34, 92–98. doi: [10.1016/j.yebeh.2014.03.016](https://doi.org/10.1016/j.yebeh.2014.03.016)
- Butcher, J. N., Hamilton, C. K., Rouse, S. V., & Cumella, E. J. (2006). The deconstruction of the Hy Scale of MMPI-2: Failure of RC3 in measuring somatic symptom expression. *Journal of Personality Assessment*, 87, 186–192. doi: [10.1207/s15327752jpa8702\\_08](https://doi.org/10.1207/s15327752jpa8702_08)
- Capilla Ramírez, P., González Ordi, H., Santamaría, P., Pérez Nieto, M. A., & Casado Morales, M. I. (2013). Fibromialgia: ¿Exageración o simulación? [Fibromyalgia: Exaggeration or malingering]. *Clínica y Salud*, 24(3), 185–195. doi: [10.5093/cl2013a20](https://doi.org/10.5093/cl2013a20)
- Carone, D. A., & Ben-Porath, Y. S. (2014). Dementia does not preclude very reliable responding on the MMP-2-RF: A case report. *The Clinical Neuropsychologist*, 28, 1019–1029. doi: [10.1080/13854046.2014.930182](https://doi.org/10.1080/13854046.2014.930182)
- Del Bene, V. A., Renteria, M. A., Maiman, M., Slugh, M., Gazzola, D.M., Nadkarni, S.S., & Barr, W.B. (2017). Increased odds and predictive rates of MMPI-2-RF scale elevations in patients with psychogenic non-epileptic seizures and observed sex differences. *Epilepsy and Behavior*, 72, 43–50. doi: [10.1016/j.yebeh.2017.04.023](https://doi.org/10.1016/j.yebeh.2017.04.023)
- Donders, J., & Pendery A (2017). Clinical Utility of the Patient Health Questionnaire-9 in the Assessment of Major Depression after Broad Spectrum Traumatic Brain Injury, *Archives of Physical Medicine and Rehabilitation*, 98, 2514–2519. doi: [10.1016/j.apmr.2017.05.019](https://doi.org/10.1016/j.apmr.2017.05.019)
- D'Orzio, L. M., Meyerowitz, B. E., Korst, L. M., Romero, R., & Goodwin, T. M. (2011). Evidence against a link between Hyperemesis Gravidarum and personality characteristics from an ethnically diverse sample of pregnant women: A pilot study. *Journal of Women's Health*, 20, 137–143. doi: [10.1089/jwh.2009.1851](https://doi.org/10.1089/jwh.2009.1851)
- Fazio, R. L., Wunderlich, W., Wilson, N., & Akeson, S. (2014). MMPI-2-RF characteristics of individuals with Interstitial Cystitis. *Journal of Psychosomatic Research*, 77, 359–362. doi: [10.1016/j.jpsychores.2014.09.010](https://doi.org/10.1016/j.jpsychores.2014.09.010)

- Forbey, J. D., Ben-Porath, Y. S., & Arbisi, P. A. (2012). The MMPI-2 Computer Adaptive version (MMPI-2-CA) in a VA medical outpatient facility. *Psychological Assessment, 24*, 628–639. doi: [10.1037/a0026509](https://doi.org/10.1037/a0026509)
- Goodpaster, K. P., Marek, R. J., Lavery, M. E., Ashton, K., Rish, J. M., & Heinberg, L. J. (2016). Graze eating among bariatric surgery candidates: Prevalence and psychosocial correlates. *Surgery for Obesity and Related Diseases, 12*, 1091-1097. doi: [10.1016/j.soard.2016.01.006](https://doi.org/10.1016/j.soard.2016.01.006)
- Granieri, A., Tamburello, S., Tamburello, A., Casale, S., Cont, C., Guglielmucci, F., & Innamorati, M. (2013). Quality of life and personality traits in patients with malignant pleural mesothelioma and their first-degree caregivers. *Neuropsychiatric Disease and Treatment, 9*, Article ID 1193–1202. doi: [10.2147/NDT.S48965](https://doi.org/10.2147/NDT.S48965)
- Heinberg, L. J., Ashton, K., Windover, A. K., & Merrell, J. (2012). Older bariatric surgery candidates: Is there greater psychological risk than for young or midlife candidates? *Surgery for Obesity and Related Diseases, 8*, 616–624. doi: [10.1016/j.soard.2011.11.005](https://doi.org/10.1016/j.soard.2011.11.005)
- Heinberg, L. J., Marek, R. J., Haskins, I. N., Hanipah, Z. N., & Brethauer, S., (in press). 30 day readmission following weight loss surgery: Can psychological factors predict non-specific indications for readmission? *Surgery for Obesity and Related Diseases*.
- Locke, D. E. C., Kirlin, K. A., Thomas, M. L., Osborne, D., Hurst, D. F., Drazkowski, J. F., Sirven, J. I., & Noe, K. H. (2010). The Minnesota Multiphasic Personality Inventory Restructured Form in the epilepsy monitoring unit. *Epilepsy and Behavior, 17*, 252–258. doi: [10.1016/j.yebeh.2009.12.004](https://doi.org/10.1016/j.yebeh.2009.12.004)
- Locke, D. E. C., Kirlin, K. A., Wershaba, R., Osborne, D., Drazkowski, J. F., Sirven, J. I., & Noe, K. H. (2011). Randomized comparison of the Personality Assessment Inventory and the Minnesota Multiphasic Personality Inventory-2, in the epilepsy monitoring unit. *Epilepsy and Behavior, 21*, 397–401. doi: [10.1016/j.yebeh.2011.05.023](https://doi.org/10.1016/j.yebeh.2011.05.023)
- Locke, D. E. C., & Thomas M. L. (2011). Initial development of Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) Scales to identify patients with psychogenic non-epileptic seizures. *Journal of Clinical and Experimental Neuropsychology, 33*, 335–343. doi: [10.1080/13803395.2010.518141](https://doi.org/10.1080/13803395.2010.518141)
- Marek, R. J., Ben-Porath, Y. S., Ashton, K., Heinberg, L. J. (2014). Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) scale score differences in bariatric surgery candidates diagnosed with binge eating disorder vs. BMI-matched controls. *International Journal of Eating Disorders, 47*, 315–319. doi: [10.1002/eat.22194](https://doi.org/10.1002/eat.22194)
- Marek, R. J., Ben-Porath, Y. S., Ashton, K., & Heinberg, L. J. (2014). Impact of using DSM-5 criteria for diagnosing binge eating disorder in bariatric surgery candidates: Change in prevalence rate, demographic characteristics, and scores on the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF). *International Journal of Eating Disorders, 47*, 553–557. doi: [10.1002/eat.22268](https://doi.org/10.1002/eat.22268)
- Marek, R. J., Ben-Porath, Y. S., Epker, J. T., Kreymer, J. K., & Block, A. R. (in press). Reliability and Validity of the Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) in Spine Surgery and Spinal Cord Stimulator Samples. *Journal of Personality Assessment*.
- Marek, R. J., Ben-Porath, Y. S., Merrell, J., Ashton, K., & Heinberg, L. J. (2014). Predicting one and three month post-operative somatic concerns, psychological distress, and maladaptive eating behavior in bariatric surgery candidates with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF). *Obesity Surgery, 24*, 631–639. doi: [10.1007/s11695-013-1149-y](https://doi.org/10.1007/s11695-013-1149-y)

- Marek, R. J., Ben-Porath, Y. S., Sellbom, M., McNulty, J. L., & Heinberg, L. J. (2015). Validity of Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) scores as a function of gender, ethnicity, and age of bariatric surgery candidates. *Surgery for Obesity and Related Diseases, 11*, 627–636. doi: [10.1016/j.soard.2014.10.005](https://doi.org/10.1016/j.soard.2014.10.005)
- Marek, R. J., Ben-Porath, Y. S., Van dulman, M., Ashton, K., & Heinberg, L. J. (2017). Using the pre-surgical psychological evaluation to predict 5-year weight-loss outcomes in bariatric surgery patients. *Surgery for Obesity and Related Diseases, 13*, 514-521. doi: [0.1016/j.soard.2016.11.008](https://doi.org/10.1016/j.soard.2016.11.008)
- Marek, R. J., Ben-Porath, Y. S., Windover, A. K., Taescavage, A. M., Merrell, J., Ashton, K., Lavery, M., & Heinberg, L. J. (2013). Assessing psychosocial functioning of bariatric surgery candidates with the Minnesota Multiphasic Personality Inventory-2-Restructured Form. *Obesity Surgery, 23*, 1864–1873. doi: [10.1007/s11695-013-1024-x](https://doi.org/10.1007/s11695-013-1024-x)
- Marek, R. J., Block, A. R., & Ben-Porath, Y. S. (2015). The Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF): Incremental validity in predicting early post-operative outcomes in spine surgery candidates. *Psychological Assessment, 27*, 114–124. doi: [10.1037/pas0000035](https://doi.org/10.1037/pas0000035)
- Marek, R. J., Block, A. R., & Ben-Porath, Y. S. (2017). Validation of a psychological screening algorithm for predicting spine surgery outcomes. *Assessment*. doi: [10.1177/1073191117719512](https://doi.org/10.1177/1073191117719512)
- Marek, R. J., Taescavage, A. M., Ben-Porath, Y. S., Ashton, K., & Heinberg, L. J. (2015). Replication and evaluation of a proposed two-factor Binge Eating Scale (BES) structure in a sample of bariatric surgery candidates. *Surgery for Obesity and Related Diseases, 11*, 659–666. doi: [0.1016/j.soard.2014.09.015](https://doi.org/10.1016/j.soard.2014.09.015)
- Marek, R. J., Taescavage, A. M., Ben-Porath, Y. S., Ashton, K., Heinberg, L. J., & Rish, J.M. (2017). Associations between psychological test results and failure to proceed with bariatric surgery. *Surgery for Obesity and Related Diseases, 13*, 507-513. doi: [10.1016/j.soard.2016.09.007](https://doi.org/10.1016/j.soard.2016.09.007)
- Marek, R. J., Taescavage, A. M., Ben-Porath, Y. S., Ashton, K., Merrell, J., Rish, J. M., & Heinberg, L. J. (2015). Using presurgical psychological testing to predict 1-year appointment adherence and weight loss in bariatric surgery candidates: Predictive validity and methodological considerations. *Surgery for Obesity and Related Diseases, 11*, 1171–1181. doi: [10.1016/j.soard.2015.03.020](https://doi.org/10.1016/j.soard.2015.03.020)
- Marek, R. J., Williams, G. A., Mohun, S. H., & Heinberg, L. J. (in press). Surgery Type and Psychosocial Factors Contribute to Poorer Weight Loss Outcomes in Persons with a Body Mass Index Greater than 60 kg/m<sup>2</sup>. *Surgery for Obesity and Related Diseases*. doi: [10.1016/j.soard.2017.09.513](https://doi.org/10.1016/j.soard.2017.09.513)
- Marinez, U., Fernandez del Rio, E., Lopez-Duran, A, Martinez-Vispo, C., & Becona, E. (2018). Types of smokers who seek smoking cessation treatment according to psychopathology. *Journal of Dual Diagnosis, 14*, 50-59. doi: [10.1080/15504263.2017.1398360](https://doi.org/10.1080/15504263.2017.1398360)
- Martinez, U., Fernandez del Rio, E., Lopez-Duran, A., & Becona, E. (2017). The utility of the MMPI-2-RF to predict the outcome of a smoking cessation treatment. *Personality and Individual Differences, 106*, 172-177. doi: [10.1016/j.paid.2016.11.019](https://doi.org/10.1016/j.paid.2016.11.019)
- McCord, D. M. & Drerup, L. C. (2011). Relative practical utility of the Minnesota Multiphasic Personality Inventory-2 Restructured Clinical Scales versus the Clinical Scales in a chronic pain patient sample. *Journal of Clinical and Experimental Neuropsychology, 33*, 140–146. doi: [10.1080/13803395.2010.495056](https://doi.org/10.1080/13803395.2010.495056)

- Merrell, J., Brethauer, S., Windover, A. K., Ashton, K., & Heinberg, L. J. (2012). Psychosocial correlates of pelvic floor disorders in women seeking bariatric surgery. *Surgery for Obesity and Related Diseases*, 8, 792–796. doi: [10.1016/j.soard.2012.01.019](https://doi.org/10.1016/j.soard.2012.01.019)
- Molvogue, M. K., Robinson, T. P., Ijaz, M. S., & McCrae, S. M. (2012). Structural neuroimaging and neuropsychological correlates of a single case of focal central pontine myelinolysis: Intact memory function with decreased intellectual and motor functions. *World Journal of Neuroscience*, 2, 192–199.
- Myers, L., Fleming, M., Lancman, M., Perrine, K., & Lancman, M. (2013). Stress coping strategies in patients with psychogenic, non-epileptic seizures and how they relate to trauma symptoms, alexithymia, anger, and mood. *Seizure*, 22, 634–639. doi: [10.1016/j.seizure.2013.04.018](https://doi.org/10.1016/j.seizure.2013.04.018)
- Myers, L., Lancman, M., Laban-Grant, O., Matzner, B., & Lancman, M. (2012). Psychogenic non-epileptic seizures: Predisposing factors to diminished quality of life. *Epilepsy and Behavior*, 25, 358–362. doi: [doi.org/10.1016/j.yebeh.2012.08.024](https://doi.org/10.1016/j.yebeh.2012.08.024)
- Myers, L., Matzner, B., Lancman, M., Perrine, K., & Lancman, M. (2013). Prevalence of alexithymia in patients with psychogenic non-epileptic seizures and epileptic seizures and predictors in psychogenic non-epileptic seizures. *Epilepsy and Behavior*, 26, 153–157. doi: [doi.org/10.1016/j.yebeh.2012.11.054](https://doi.org/10.1016/j.yebeh.2012.11.054)
- Myers, L., Perrine, K., Lancman, M., Fleming, M., & Lancman, M. (2013). Psychological trauma in patients with psychogenic non-epileptic seizures: Trauma characteristics and those who develop PTSD. *Epilepsy and Behavior*, 28, 121–126. doi: [10.1016/j.yebeh.2013.03.033](https://doi.org/10.1016/j.yebeh.2013.03.033)
- Nelson, N. W., Hoelzle, J. B., McHuire, K. A., Sim, A. H., Goldman, D. J., Ferrier-Aurbach, A. G., Charlesworth, M. J., Arbisi, P. A., & Sponheim, S. R. (2011). Self-report of psychological function among OEF/OIF personnel who also report combat-related concussion. *The Clinical Neuropsychologist*, 25, 716–740. doi: [10.1080/13854046.2011.579174](https://doi.org/10.1080/13854046.2011.579174)
- Pona, A. A., Heinberg, L. J., Lavery, M., Ben-Porath, Y. S., & Rish, J. M., (2016). Psychological predictors of body image concerns 3-months after bariatric surgery. *Surgery for Obesity and Related Diseases*, 12, 188-193. doi: [10.1016/j.soard.2015.05.008](https://doi.org/10.1016/j.soard.2015.05.008)
- Pona, A. A., Marek, R. J., Heinberg, L. J., Lavery, M., Ashton, K., & Rish, J. M. (2017). Psychological correlates of body image dissatisfaction before and after bariatric surgery. *Bariatric Surgical Practice and Patient Care*. doi: [10.1089/bari.2017.0021](https://doi.org/10.1089/bari.2017.0021)
- Stroescu, I., Salinas, C. M., Nahab, F. B., & Stringer, A. Y. (2011). Long-term neurocognitive and neuroimaging outcomes in posterior reversible Encephalopathy Syndrome: Two case reports and implications. *The Clinical Neuropsychologist*, 25, 1386–1402. doi: [10.1080/13854046.2011.628339](https://doi.org/10.1080/13854046.2011.628339)
- Tarescavage, A. M., Scheman, J., & Ben-Porath, Y. S. (2015). Reliability and validity of the Minneapolis Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) in evaluations of chronic low back pain patients. *Psychological Assessment*, 27, 433–446. doi: [10.1037/pas0000056](https://doi.org/10.1037/pas0000056)
- Tarescavage, A. M., Scheman, J., & Ben-Porath, Y. S. (2018). Prospective comparison of the Minnesota Multiphasic Personality Inventory-2 (MMPI-2) and MMPI-2 Restructured Form (MMPI-2-RF) in predicting treatment outcomes among patients with chronic low back pain. *Journal of Clinical Psychology in Medical Settings*, 25, 66-79. doi: [10.1007/s10880-017-9535-6](https://doi.org/10.1007/s10880-017-9535-6)

- Tarescavage, A. M., Windover, A. K., Ben-Porath, Y. S., Botuacoff, L. I., Marek, R. J., Ashton, K., Merrell, J., Lavery, M., & Heinberg, L. J. (2013). Use of the MMPI-2-RF Suicidal/Death Ideation and Substance Abuse Scales in screening bariatric surgery candidates. *Psychological Assessment, 25*, 1384–1389. doi: [10.1037/a0034045](https://doi.org/10.1037/a0034045)
- Tarescavage, A. M., Wygant, D. B., Boutacoff, L. I., & Ben-Porath, Y. S. (2013). Reliability, validity, and utility of the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) in assessments of bariatric surgery candidates. *Psychological Assessment, 25*, 1179–1194. doi: [10.1037/a0033694](https://doi.org/10.1037/a0033694)
- Thomas, M. L., & Locke, D. E. C. (2010). Psychometric properties of the MMPI-2-RF Somatic Complaints (RC1) Scale. *Psychological Assessment, 22*, 492–503. doi: [10.1037/a0019229](https://doi.org/10.1037/a0019229)
- Walvoort, S. J. W., van der Heijden, P. T., Wester, A. J., Kessels, R. P. C., & Egger, J. I. M. (2016). Self-awareness of cognitive dysfunction: Self-reported complaints and cognitive performance in patients with alcohol-induced mild or major neurocognitive disorder. *Psychiatry Research, 245*, 291–296. doi: [10.1016/j.psychres.2016.08.007](https://doi.org/10.1016/j.psychres.2016.08.007)
- Wygant, D. B., Boutacoff, L. A., Arbisi, P. A., Ben-Porath, Y. S., Kelly, P. H., & Rupp, W. M. (2007). Examination of the MMPI-2 Restructured Clinical (RC) Scales in a sample of bariatric surgery candidates. *Journal of Clinical Psychology in Medical Settings, 14*, 197–205. doi: [10.1007/s10880-007-9073-8](https://doi.org/10.1007/s10880-007-9073-8)
- Yamout, K. Z., Heimrichs, R. J., Baade, L. E., Soetaert, D. K., & Liow, K. K. (2017). Comparative prediction of nonepileptic events using MMPI-2 clinical scales Harris Lingoes subscales, and restructured clinical scales. *Epilepsy & Behavior, 68*, 31–34. doi: [10.1016/j.yebeh.2016.12.008](https://doi.org/10.1016/j.yebeh.2016.12.008)

### **MENTAL HEALTH SETTINGS**

- Anderson, J. L., Sellbom, M., Ayearst, L., Quilty, L. C., Chmielewski, M., & Bagby, R. M. (2015). Associations between *DSM-5* Section III personality traits and the Minnesota Multiphasic Personality Inventory 2-Restructured Form (MMPI-2-RF) scales in a psychiatric patient sample. *Psychological Assessment, 27*, 811–815. doi: [10.1037/pas0000096](https://doi.org/10.1037/pas0000096)
- Anderson, J. L., Sellbom, M., Sansone, R. A., & Songer, D. A. (2016). Comparing external correlates of *DSM-5* Section II and Section III borderline personality disorder constructs. *Journal of Personality Disorders, 30*, 193–210. doi: [10.1521/pepi\\_2015\\_29\\_189](https://doi.org/10.1521/pepi_2015_29_189)
- Anestis, J. C., Finn, J. A., Gottfried, E. D., Arbisi, P. A., & Joiner, T. E. (2015). Reading the road signs: The utility of the MMPI-2 Restructured Form Validity Scales in prediction of premature termination. *Assessment, 22*, 279–288. doi: [073191114541672](https://doi.org/073191114541672)
- Anestis, J. C., Gottfried, E. D., & Joiner, T. E. (2015). The utility of MMPI-2-RF substantive scales in prediction of negative treatment outcomes in a community mental health center. *Assessment, 22*, 23–35. doi: [10.1177/1073191114536771](https://doi.org/10.1177/1073191114536771)
- Anestis, J. C., Finn, J. A., Gottfried, E. D., Hames, J. L., Bodell, L. P., Hagan, C. R., Arnau, R. C., Anestis, M. D., Arbisi, P. A., & Joiner, T. E. (2018). Burdonesomeness, Belongingness, and capability: Assessing the interpersonal-psychological theory of suicide with MMPI-2-RF scales. *Assessment, 25*, 415–431. doi: [10.1177/1073191116652227](https://doi.org/10.1177/1073191116652227)
- Arbisi, P. A., Erbes, C. R., Polusny, M. A., & Nelson, N. W. (2010). The concurrent and incremental validity of the Trauma Symptom Inventory in women reporting histories of sexual maltreatment. *Assessment, 17*, 406–418. doi: [10.1177/1073191110376163](https://doi.org/10.1177/1073191110376163)



- Arbisi, P. A., Polusny, M. A., Erbes, C. R., Thusras, P., & Reddy, M. K. (2011). The Minnesota Multiphasic Personality Inventory-2 Restructured Form in National Guard soldiers screening positive for posttraumatic stress disorder and mild traumatic brain injury. *Psychological Assessment, 23*, 203–214. doi: [10.1037/a0021339](https://doi.org/10.1037/a0021339)
- Arbisi, P. A., Rusch, L., Polusny, M. A., Thuras, P., & Erbes, C. R. (2013). Does cynicism play a role in failure to obtain needed care? Mental health service utilization among returning U.S. National Guard soldiers. *Psychological Assessment, 25*, 991–996. doi: [10.1037/a0032225](https://doi.org/10.1037/a0032225)
- Arbisi, P. A., Sellbom, M., & Ben-Porath, Y. S. (2008). Empirical correlates of the MMPI-2 Restructured Clinical (RC) Scales in psychiatric inpatients. *Journal of Personality Assessment, 90*, 122–128. doi: [10.1080/00223890701845146](https://doi.org/10.1080/00223890701845146)
- Bagby, R. M., Sellbom, M., Ayearst, L. E., Chmielewski, M. S., Anderson, J. L., & Quilty, L. C. (2014). Exploring the Hierarchical Structure of the MMPI-2-RF Personality Psychopathology Five in Psychiatric Patient and University Student Samples. *Journal of Personality Assessment, 96*, 166–172. doi: [10.1080/00223891.2013.825623](https://doi.org/10.1080/00223891.2013.825623)
- Ben-Porath, Y. S., & Tellegen, A. (2018). Leone, Mosticoni, Biondi, and Butcher's (2018) effort to compare the MMPI-2-RF with the MMPI-2 falls well short. *Archives of Assessment Psychology, 8*, 23–31.
- Binford, A., & Liljequist, L. (2008). Behavioral correlates of selected MMPI-2 Clinical, Content, and Restructured Clinical Scales. *Journal of Personality Assessment, 60*, 608–614. doi: [10.1080/00223890802388657](https://doi.org/10.1080/00223890802388657)
- Bolinsky, P. K., Trumbetta, S. L., Hanson, D. R., & Gottesman, I. I. (2010). Predicting adult psychopathology from adolescent MMPIs: Some victories. *Personality and Individual Differences, 49*, 324–330. doi: [10.1016/j.paid.2010.01.026](https://doi.org/10.1016/j.paid.2010.01.026)
- Bosch, P., Van Luijtelaar, G., Van Den Noort, M., Schenkwald, J., Kueppenbender, N., Lim, S., Egger, J., & Coenen, A. (2014). The MMPI-2 in chronic psychiatric illness. *Scandinavian Journal of Psychology, 55*, 513–519. doi: [10.1111/sjop.12152](https://doi.org/10.1111/sjop.12152)
- Bryant, W. T., & McNulty, J. L. (2017). Which domain of the PSY-5 is most relevant to substance use? *Journal of Personality Assessment, 99*, 524–533. doi: [10.1080/00223891.2016.1250213](https://doi.org/10.1080/00223891.2016.1250213)
- Castro, Y., Gordon, K. H., Brown, J. S., Cox, J. C., & Joiner, T. E. (2008). Examination of racial differences on the MMPI-2 Clinical and Restructured Clinical Scales in an outpatient sample. *Assessment, 15*, 277–286. doi: [10.1177/1073191107312735](https://doi.org/10.1177/1073191107312735)
- Clark, R., DeYoung, C. G., Sponheim, S. R., Bender, T. L., Polusny, M. A., Erbes, C. R., & Arbisi, P. A. (2013). Predicting post-traumatic stress disorder in veterans: Interaction of traumatic load with COMT gene variation. *Journal of Psychiatric Research, 47*, 1849–1856. doi: [10.1007/s10862-013-9358-1](https://doi.org/10.1007/s10862-013-9358-1)
- Choi, J. Y. (2017). Posttraumatic stress symptoms and dissociation between childhood trauma and two different types of psychosis-like experience. *Child Abuse and Neglect, 72*, 404–410. doi: [10.1016/j.chiabu.2017.08.023](https://doi.org/10.1016/j.chiabu.2017.08.023)
- Durosini, I., Tarocchi, A., & Aschieri, F. (2017). Therapeutic assessment with a client with persistent complex bereavement disorder: A single-case time-series design. *Clinical Case Studies*. doi: [10.1177/1534650117693942](https://doi.org/10.1177/1534650117693942)
- Erbes, C. R., Kramer, M., Arbisi, P. A., DeGarmo, D., & Polusny, M. A. (2017). Characterizing spouse/partner depression and alcohol problems of the course of military deployment. *Journal of Consulting and Clinical Psychology, 85*, 297–308. doi: [10.1037/ccp0000190](https://doi.org/10.1037/ccp0000190)

- Erbes, C. R., Polusny, M. A., Arbisi, P. A., & Koffel, E. (2012). PTSD symptoms in a cohort of National Guard Soldiers deployed to Iraq: Evidence for nonspecific and specific components. *Journal of Affective Disorders, 142*, 269–274. doi: [10.1016/j.jad.2012.05.013](https://doi.org/10.1016/j.jad.2012.05.013)
- Finn, J. A., Arbisi, P. A., Erbes, C. R., Polusny, M. A., & Thuras, P. (2014). The MMPI-2 Restructured Form Personality Psychopathology Five Scales: Bridging DSM-5 Section 2 personality disorders and DSM-5 Section 3 personality trait dimensions. *Journal of Personality Assessment, 96*, 173–184. doi: [00223891.2013.866569](https://doi.org/00223891.2013.866569)
- Forbes, D., Elhai, J. D., Miller, M. W., & Creamer, M. (2010). Internalizing and externalizing classes in posttraumatic stress disorder: A latent class analysis. *Journal of Traumatic Stress, 23*, 340–349. doi: [10.1002/jts.20526](https://doi.org/10.1002/jts.20526)
- Gilbert, S. E., & Gordon, K. C. (2013). Interpersonal psychotherapy informed treatment for avoidant personality disorder with subsequent depression. *Clinical Case Studies, 12*, 111–127. doi: [10.1177/1534650112468611](https://doi.org/10.1177/1534650112468611)
- Gordon, R. M., Stoffey, R. W., & Perkins, B. L. (2013). Comparing the sensitivity of the Clinical Scales and the MMPI-RC (sic) Scales to clients rated as psychotic, borderline, or neurotic on the Psychodiagnostic Chart. *Psychology, 4*, 12–16. doi: [10.4236/psych.2013.49A1003](https://doi.org/10.4236/psych.2013.49A1003)
- Gottfried, E., Bodell, L. M., Carbonell, J., & Joiner, T. (2014). The clinical utility of the MMPI-2-RF Suicidal/Death Ideation Scale. *Psychological Assessment, 26*, 1205–1211. doi: [10.1037/pas0000017](https://doi.org/10.1037/pas0000017)
- Haber, J. C., & Baum, L. J. (2014). Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) Scales as predictors of psychiatric diagnoses. *South African Journal of Psychology, 44*, 439–453. doi: [10.1177/0081246314532788](https://doi.org/10.1177/0081246314532788)
- Handel, R. W., & Archer, R. P. (2008). An investigation of the psychometric properties of the MMPI-2 Restructured Clinical (RC) Scales with mental health inpatients. *Journal of Personality Assessment, 90*, 239–249. doi: [10.1080/00223890701884954](https://doi.org/10.1080/00223890701884954)
- Kamphuis, J. H., Arbisi, P. A., Ben-Porath, Y. S., & McNulty, J. L. (2008). Detecting comorbid Axis-II status among inpatients using the MMPI-2 Restructured Clinical Scales. *European Journal of Psychological Assessment, 24*, 157–164. doi: [10.1027/1015-5759.24.3.157](https://doi.org/10.1027/1015-5759.24.3.157)
- Koffel, E., Kramer, M. D., Arbisi, P. A., Erbes, C. R., Kaler, M., & Polusny, M. A. (2015). Personality traits and combat exposure as predictors of psychopathology over time. *Psychological Medicine*. doi: [10.1017/S0033291715001798](https://doi.org/10.1017/S0033291715001798)
- Koffel, E., Polusny, M. A., Arbisi, P. A., & Erbes, C. R. (2012). A preliminary investigation of the new and revised symptoms of posttraumatic stress disorder in DSM-5. *Depression and Anxiety, 29*, 731–73. doi: [10.1002/da.21965](https://doi.org/10.1002/da.21965)
- Lee, Y. T. C., Graham, J. R., & Arbisi, P. A. (2018). The utility of MMPI-2-RF scale scores in differential diagnosis of Schizophrenia and Major Depressive Disorder. *Journal of Personality Assessment, 100*, 305–312. doi: [10.1080/00223891.2017.1300906](https://doi.org/10.1080/00223891.2017.1300906)
- Leone, C., Mosticoni, S., Iannella, F., Biondi, M., & Butcher, J. N. (2018). Comparability of MMP-2-RF with MMPI-2 in assessing psychiatric patients: A shortfall. *Archives of Assessment Psychology, 8*, 7–22.
- McDevitt-Murphy, M. E., Weathers, F. W., Flood, A. M., Eakin, D. E., & Benson, T. A. (2007). The utility of the PAI and the MMPI-2 for discriminating PTSD, depression, and social phobia in trauma-exposed college students. *Assessment, 14*, 181–195. doi: [10.1177/1073191106295914](https://doi.org/10.1177/1073191106295914)

- McManus, E. S., Cuccurullo, L. J., Uddo, M., & Franklin, C. L. (2018). MMPI-2-RF characteristics of veterans seeking treatment for military sexual trauma. *Psychological Assessment, 30*, 561-266. doi: [10.1037/pas0000527](https://doi.org/10.1037/pas0000527)
- Meyers, J. E., Grills, C. E., Zellinger, M. M., Miller, R. M. (2014). Emotional distress affects attention and concentration: The difference between mountains and valleys. *Applied Neuropsychology: Adult, 21*, 28-35. doi: [10.1080/09084282.2012.721148](https://doi.org/10.1080/09084282.2012.721148)
- Meyers, J. E., Miller, R. M., & Tuita, A. R. R. (2014). Using pattern analysis matching to differentiate TBI and PTSD in a military sample. *Applied Neuropsychology: Adult, 21*, 60-68. doi: [10.1080/09084282.2012.737881](https://doi.org/10.1080/09084282.2012.737881)
- Miller, S. N., Bozzay, M. L., Ben-Porath, Y. S., & Arbisi, P. A. (2017). Distinguishing levels of suicide risk in depressed male veterans: The role of internalizing and externalizing psychopathology as measured by the MMPI-2-RF. *Assessment*. doi: [10.1177/1073191117743787](https://doi.org/10.1177/1073191117743787)
- Miller, M. W., Wolf, E. J., Harrington, K. M., Brown, T. A., Kaloupek, D. G., & Keane, T. M. (2010). An evaluation of competing models for the structure of PTSD symptoms using external measures of comorbidity. *Journal of Traumatic Stress, 23*, 631-638. doi: [10.1002/jts.20559](https://doi.org/10.1002/jts.20559)
- Moutrie, J. K., & Engel, R. R. (2016). Empirical correlates for the Minnesota Multiphasic Personality Inventory-2-Restructured Form in a German inpatient sample. *Psychological Assessment*. doi: [10.1037/pas0000415](https://doi.org/10.1037/pas0000415)
- Noordhof, A., Sellbom, M., Eigenhuis, A., & Kamphuis, J. H. (2015). Distinguishing between demoralization and specific personality traits in clinical assessment with the NEO-PI-R. *Psychological Assessment, 27*, 645-655. doi: [10.1037/pas0000067](https://doi.org/10.1037/pas0000067)
- Ozonoff, S., Garcia, N., Clark, E., & Lainhart, J. E. (2005). MMPI-2 personality profiles of high-functioning adults with autism spectrum disorders. *Assessment, 12*, 86-95. doi: [10.1177/1073191104273132](https://doi.org/10.1177/1073191104273132)
- Purdon, S. E., Purser, S. M., & Goddard, K. M. (2011). MMPI-2 Restructured Form over-reporting scales in first-episode psychosis. *The Clinical Neuropsychologist, 25*, 829-842. doi: [10.1080/13854046.2011.585141](https://doi.org/10.1080/13854046.2011.585141)
- Quilty, L. C., Sellbom, M., Tackett, J. L., & Bagby, R. M. (2009). Personality trait predictors of bipolar disorder: A replication and extension in a psychiatric sample. *Psychiatry Research, 169*, 159-163. doi: [10.1016/j.psychres.2008.07.004](https://doi.org/10.1016/j.psychres.2008.07.004)
- Reid, R. C., & Carpenter, B. N. (2009). Demoralization, hypomanic activation, and disinhibition scores on MMPI-2 as significant predictors of hypersexual behavior. *Sexual Addiction and Compulsivity, 16*, 173-189. doi: [10.1080/10720160903202448](https://doi.org/10.1080/10720160903202448)
- Reid, R. C., & Carpenter, B. N. (2009). Exploring relationships of psychopathology in hypersexual patients using the MMPI-2. *Journal of Sex & Marital Therapy, 35*(4), 294-310. doi: [10.1080/00926230902851298](https://doi.org/10.1080/00926230902851298)
- Reid, R. C., Carpenter, B. N., & Draper, E. D. (2011). Disputing the notion of psychopathology among women married to hypersexual men using the MMPI-2-RF. *Journal of Sex & Marital Therapy, 37*(1), 45-55. doi: [10.1080/0092623X.2011.533585](https://doi.org/10.1080/0092623X.2011.533585)
- Rogers, M. L., Anestis, J. C., Harrop, T. M., Schneider, M., Bender, T. W., Ringer, F. B., & Joiner, T. E. (2017). Examination of MMPI-2-RF substantive scales as indicators of acute suicidal affective disturbance components. *Journal of Personality Assessment, 99*, 424-434. doi: [10.1080/00223891.2016.1222393](https://doi.org/10.1080/00223891.2016.1222393)

- Scholte, W., Tiemens, B. G., Verheul, R., Meerman, A., Eggers, J., & Hutschemaekers, G. (2012). Predictive validity of the MMPI-2 Clinical, PSY-5, and RC Scales for therapy disruptive behavior. *Journal of Psychiatric Practice, 18*, 420–429. doi: [10.1097/01.pra.0000422740.87495.91](https://doi.org/10.1097/01.pra.0000422740.87495.91)
- Scholte, W., Tiemens, B. G., Verheul, R., Meerman, A., Eggers, J., & Hutschemaekers, G. (2012). The RC Scales predict psychotherapy outcomes: The predictive validity of the MMPI-2's Restructured Clinical Scales for psychotherapeutic outcomes. *Personality and Mental Health, 6*, 292–302. doi: [10.1002/pmh.1190](https://doi.org/10.1002/pmh.1190)
- Sellbom, M., Bagby, R. M., Kushner, S., Quilty, L. C., & Ayearst, L. E. (2012). Diagnostic construct validity of the MMPI-2 Restructured Form (MMPI-2-RF) scale scores. *Assessment, 19*, 176–186. doi: [10.1177/1073191111428763](https://doi.org/10.1177/1073191111428763)
- Sellbom, M., Ben-Porath, Y. S., & Bagby, R. M. (2008). Personality and psychopathology: Mapping the MMPI-2 Restructured Clinical (RC) Scales onto the Five Factor Model of Personality. *Journal of Personality Disorders, 22*, 291–312. doi: [10.1521/pedi.2008.22.3.291](https://doi.org/10.1521/pedi.2008.22.3.291)
- Sellbom, M., Ben-Porath, Y. S., & Graham, J. R. (2006). Correlates of the MMPI-2 Restructured Clinical (RC) Scales in a college counseling setting. *Journal of Personality Assessment, 86*, 89–99. doi: [10.1207/s15327752jpa8601\\_10](https://doi.org/10.1207/s15327752jpa8601_10)
- Sellbom, M., Ben-Porath, Y. S., McNulty, J. L., Arbisi, P. A., & Graham, J. R. (2006). Elevation differences between MMPI-2 Clinical and Restructured Clinical (RC) Scales: Frequency, origins, and interpretative implications. *Assessment, 13*, 430–441. doi: [10.1177/1073191106293349](https://doi.org/10.1177/1073191106293349)
- Sellbom, M., Graham, J. R., & Schenk, P. (2006). Incremental validity of the MMPI-2 Restructured Clinical (RC) Scales in a private practice sample. *Journal of Personality Assessment, 86*, 196–205. doi: [10.1207/s15327752jpa8602\\_09](https://doi.org/10.1207/s15327752jpa8602_09)
- Sellbom, M., Waugh, M. H., & Hopwood, C. J. (2018). Development and Validation of Personality Disorder Spectra Scales for the MMPI-2-RF. *Journal of Personality Assessment*. doi: [10.1080/00223891.2017.1407327](https://doi.org/10.1080/00223891.2017.1407327)
- Shkalim, E. (2015). Psychometric evaluation of the MMPI-2/MMPI-2-RF Restructured Clinical Scales in an Israeli sample. *Assessment, 22*, 607–618. doi: [10.1177/1073191114555884](https://doi.org/10.1177/1073191114555884)
- Shklalim, E., Almagor, M., & Ben-Porath, Y. S. (2017). Examining current conceptualizations of psychopathology with the MMPI-2/MMPI-2-RF Restructured Clinical Scales: Preliminary findings from a cross-cultural study. *Journal of Personality Assessment, 99*, 375–383. doi: [10.1080/00223891.2016.1189429](https://doi.org/10.1080/00223891.2016.1189429)
- Shkalim, E., Ben-Porath, Y. S., & Almagor, M. (2016). Mapping the MMPI-2/MMPI-2-RF Restructured Clinical Scales onto mood markers in an Israeli sample. *Journal of Personality Assessment, 98*, 430–434. doi: [10.1080/00223891.2016.1189429](https://doi.org/10.1080/00223891.2016.1189429)
- Simms, L. J., Casillas, A., Clark, L. A., Watson, D., & Doebbeling, B. I. (2005). Psychometric evaluation of the Restructured Clinical Scales of the MMPI-2. *Psychological Assessment, 17*, 345–358. doi: [10.1037/1040-3590.17.3.345](https://doi.org/10.1037/1040-3590.17.3.345)
- Suhr, J. A., Cook, C., & Morgan, B. (2017). Assessing functional impairment in ADHD: Concerns for validity of self-report. *Psychological Injury and Law, 10*, 151–170. doi: [10.1007/s12207-017-9292-8](https://doi.org/10.1007/s12207-017-9292-8)

- Tarescavage, A. M., Finn, J. A., Marek, R. J., Ben-Porath, Y. S., & van Dulmen, M. H. M. (2015). Premature termination from psychotherapy and internalizing psychopathology: The role of demoralization. *Journal of Affective Disorders, 174*, 549–555. doi: [10.1016/j.jad.2014.12.018](https://doi.org/10.1016/j.jad.2014.12.018)
- Vachon, D. D., Sellbom, M., Ryder, A. G., Miller, J. D., & Bagby, R. M. (2009). A Five-Factor Model description of depressive personality disorder. *Journal of Personality Disorders, 23*, 447–465. doi: [10.1521/pepi.2009.23.5.447](https://doi.org/10.1521/pepi.2009.23.5.447)
- van der Heijden, P. T., Egger, J. I. M., Rossi, G., & Derksen, J. J. L. (2012). Integrating psychopathology and personality disorders conceptualized by the MMPI-2-RF and the MCMI-III: A structural validity study. *Journal of Personality Assessment, 94*, 345–347. doi: [10.1080/00223891.2012.656861](https://doi.org/10.1080/00223891.2012.656861)
- van der Heijden, P. T., Egger, J. I. M., Rossi, G., & Derksen, J. J. L. (2013). The MMPI-2-Restructured Form and the standard MMPI-2 Clinical Scales in relation to *DSM-IV*. *European Journal of Psychological Assessment, 29*, 182–188. doi: [10.1027/1015-5759/a000140](https://doi.org/10.1027/1015-5759/a000140)
- van der Sterren-Kusters, W. J. C., van der Heijden, P. T., & Egger, J. I. M. (2017). Psychometric properties of the Dutch Anxiety Change Expectancy Scale (ACES-NL). *International Journal of Psychology and Psychological Therapy, 17*, 189–198.
- Venables, N. C., Sellbom, M., Sourander, A., Kendler, K. S., Joiner, T. E., Drislane, L. E., Sillanmaki, L., Elonheimo, H., Parkola, K., Multimaki, P., & Patrick, C. J. (2015). Separate and interactive contributions of weak inhibitory control and threat sensitivity to prediction of suicide risk. *Psychiatry Research, 226*, 461–466. doi: [10.1016/j.psychres.2015.01.018](https://doi.org/10.1016/j.psychres.2015.01.018)
- Wallace, A., & Liljequist, L. (2005). A comparison of the correlational structures and elevation patterns of the MMPI-2 Restructured Clinical (RC) and Clinical Scales. *Assessment, 12*, 290–294. doi: [10.1177/1073191105276250](https://doi.org/10.1177/1073191105276250)
- Watson, C., Quilty, L. C., & Bagby, R. M. (2011). Differentiating bipolar disorder from major depressive disorder using the MMPI-2-RF: A receiver operating characteristics (ROC) analysis. *Journal of Psychopathology and Behavioral Assessment, 33*, 368–374. doi: [10.1007/s10862-010-9212-7](https://doi.org/10.1007/s10862-010-9212-7)
- Wolf, E. J., Miller, M. W., Orazem, R. J., Weierich, M. R., Castillo, D. T., Milford, J., Kaloupek, D. G., & Keane, T. M. (2008). The MMPI-2 Restructured Clinical Scales in the assessment of posttraumatic stress disorder and comorbid disorders. *Psychological Assessment, 20*, 327–340. doi: [10.1037/a0012948](https://doi.org/10.1037/a0012948)
- Wygant, D. B., & Fleming, K. P. (2008). Clinical utility of the MMPI-2 Restructured Clinical (RC) Scales in therapeutic assessment: A case study. *Journal of Personality Assessment, 90*, 110–118. doi: [10.1080/00223890701845112](https://doi.org/10.1080/00223890701845112)
- Zahn, N., Sellbom, M., Pymont, C., & Schenk, P.W. (2017). Associations Between MMPI-2-RF Scale Scores and Self-Reported Personality Disorder Criteria in a Private Practice Sample. *Journal of Psychopathology and Behavioral Assessment*. doi: [10.1007/s10862-017-9616-8](https://doi.org/10.1007/s10862-017-9616-8)

## **ADDICTION AND SUBSTANCE ABUSE TREATMENT SETTINGS**

- Forbey, J. D., & Ben-Porath, Y. S. (2007). A comparison of the MMPI-2 Restructured Clinical (RC) and Clinical Scales in a substance abuse treatment sample. *Psychological Services, 4*, 46–58. doi: [10.1037/1541-1559.4.1.46](https://doi.org/10.1037/1541-1559.4.1.46)



- Hopkins, T. A., Brawner, C. A., Meyer, M., Zawalinski, L., Carnes, P. J., & Green, B. A. (2016). MMPI-2 correlates of sadomasochism in a sexual addiction sample: Contrasting men and women. *Sexual Addiction & Compulsivity, 23*, 114-140. doi: [10.1080/10720162.2015.1095137](https://doi.org/10.1080/10720162.2015.1095137)
- Monnot, M. J., Quirk, S. W., Hoerger, M., & Brewer, L. (2009). Racial bias in personality assessment: Using the MMPI-2 to predict psychiatric diagnoses of African American and Caucasian chemical dependency patients. *Psychological Assessment, 21*, 137-151. doi: [10.1037/a0015316](https://doi.org/10.1037/a0015316)

## **POLICE AND PUBLIC SAFETY SETTINGS**

- Corey, D. M., Sellbom, M., & Ben-Porath, Y. S., (in press). Risks associated with overcontrolled behavior in police officer recruits. *Psychological Assessment*.
- Detrick, P., Ben-Porath, Y. S., & Sellbom, M. (2015). Associations between MMPI-2-RF (Restructured Form) and Inwald Personality Inventory (IPI) scale scores in a law enforcement preemployment screening sample. *Journal of Police and Criminal Psychology*. doi: [10.1007/s11896-015-9172-7](https://doi.org/10.1007/s11896-015-9172-7)
- Detrick, P., & Chibnall, J. T. (2014). Underreporting on the MMPI-2-RF in a high demand police officer selection content: An illustration. *Psychological Assessment, 26*, 1044-1049. doi: [10.1037/pas0000013](https://doi.org/10.1037/pas0000013)
- Roberts, R. M., Tarescavage, A. M., Ben-Porath, Y. S., & Roberts, M. D. (2018). Predicting Post-Probationary Job Performance of Police Officers Using CPI and MMPI-2 RF Test Data Obtained During Preemployment Psychological Screening. *Journal of Personality Assessment*. doi: [1080/00223891.2018.1423990](https://doi.org/1080/00223891.2018.1423990)
- Sellbom, M., Fischler, G. L., & Ben-Porath, Y. S. (2007). Identifying MMPI-2 predictors of police officer integrity and misconduct. *Criminal Justice and Behavior, 34*, 985-1004. doi: [10.1177/0093854807301224](https://doi.org/10.1177/0093854807301224)
- Tarescavage, A. M., Brewster, J., Corey, D. M., & Ben-Porath, Y. S. (2015). Use of pre-hire MMPI-2-RF police candidate scores to predict supervisor ratings of post-hire performance. *Assessment*. doi: [10.1177/1073191114548445](https://doi.org/10.1177/1073191114548445)
- Tarescavage, A. M., Corey, D. M., & Ben-Porath, Y. S. (2015). Minnesota Multiphasic Personality Inventory Restructured Form (MMPI-2-RF) predictors of police officer problem behavior. *Assessment, 22*(1), 116-132. doi: [10.1177/1073191114534885](https://doi.org/10.1177/1073191114534885)
- Tarescavage, A. M., Corey, D. M., & Ben-Porath, Y. S. (2015). A prorating method for estimating MMPI-2-RF Scores from MMPI responses: Examination of score fidelity and illustration of empirical utility in the PERSEREC Police Integrity Study sample. *Assessment*. doi: [10.1177/1073191115575070](https://doi.org/10.1177/1073191115575070)
- Tarescavage, A. M., Corey, D. M., Gupton, H. M., & Ben-Porath, Y. S. (2015). Criterion validity and clinical utility of the Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) in assessments of police officer candidates. *Journal of Personality Assessment*. doi: [10.1080/00223891.2014.995800](https://doi.org/10.1080/00223891.2014.995800)
- Tarescavage, A. M., Fischler, G. L., Cappo, B., Hill, D. O., Corey, D. M., & Ben-Porath, Y. S. (2015). Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) predictors of police officer problem behavior and collateral self-report test scores. *Psychological Assessment, 27*, 125-137. doi: [10.1037/pas0000041](https://doi.org/10.1037/pas0000041)

## **OTHER NON-CLINICAL SETTINGS**

- Anderson, J. L., Sellbom, M., & Salekin, R. T. (2018). Utility of the Personality Inventory for DSM-5 Brief Form (PID-5-BF) in the measurement of Section II personality disorders and maladaptive behaviors. *Assessment, 25*, 596-607. doi: [10.1177/1073191116676889](https://doi.org/10.1177/1073191116676889)

- Anderson, J. L., Sellbom, M., Sansone, R. A., & Songer, D. A. (2015). Comparing external correlates of DSM-5 Section II and Section III dimensional trait operationalizations of Borderline Personality Disorder. *Journal of Personality Disorders*. doi: [10.1521/pepi.2015.29.189](https://doi.org/10.1521/pepi.2015.29.189)
- Avdeyeva, T. V., Tellegen, A., & Ben-Porath, Y. S. (2011). Empirical correlates of low scores on MMPI-2/MMPI-2-RF Restructured Clinical Scales in a sample of university students. *Assessment*, *19*, 388–393. doi: [10.1177/10731911111411675](https://doi.org/10.1177/10731911111411675)
- Ayearst, L. E., Sellbom, M., Trobst, K. K., & Bagby, R. M. (2013). Evaluating the interpersonal content of the MMPI-2-RF Interpersonal Scales. *Journal of Personality Assessment*, *95*, 187–196. doi: [10.1080/00223891.2012.730085](https://doi.org/10.1080/00223891.2012.730085)
- Baker, C. N., & Hoerger, M. (2012). Parental child-rearing strategies influence self-regulation, socio-emotional adjustment, and psychopathology in early adulthood: Evidence from a retrospective cohort study. *Personality and Individual Differences*, *52*, 800–805. doi: [10.1016/j.paid.2011.12.034](https://doi.org/10.1016/j.paid.2011.12.034)
- Benitez, A., & Gunstad, J. (2012). Poor sleep quality diminishes cognitive functioning independent of depression and anxiety in healthy young adults. *The Clinical Neuropsychologist*, *26*, 214–223. doi: [10.1080/13854046.2012.658439](https://doi.org/10.1080/13854046.2012.658439)
- Bolinski, P. K., Guidi, J. P., Myers, K. R., Cooper-Bolinski, D., Sschuder, K. M., James, A. V., Hudak, D. V., Gonzales, Y., McTierman, E., & Smith, E. (2016). The MMPI-2-RF and college students: Do we remain stuck in a normative no-man's land? *Archives of Assessment Psychology*, *6*, 81–95.
- Brinker, J. K., Chin, Z. H., & Wilkinson, R. (2014). Ruminative thinking style and the MMPI-2-RF. *Personality and Individual Differences*, *66*, 102–105. doi: [10.1016/j.paid.2014.03.001](https://doi.org/10.1016/j.paid.2014.03.001)
- Forbey, J. D., & Ben-Porath, Y. S. (2008). Empirical correlates of the MMPI-2 Restructured Clinical (RC) Scales in a non-clinical setting. *Journal of Personality Assessment*, *90*, 136–141. doi: [10.1080/00223890701845161](https://doi.org/10.1080/00223890701845161)
- Forbey, J. D., & Lee, T. T. C. (2011). An exploration of the impact of invalid MMPI-2 protocols on collateral self-report measure scores. *Journal of Personality Assessment*, *93*, 556–565. doi: [10.1080/00223891.2011.608757](https://doi.org/10.1080/00223891.2011.608757)
- Forbey, J. D., Lee, T. T. C., & Handel, R. W. (2010). Correlates of the MMPI-2-RF in a college setting. *Psychological Assessment*, *22*, 737–744. doi: [10.1037/a0020645](https://doi.org/10.1037/a0020645)
- Franz, A. O., Harrop, T. M., & McCord, D. M. (2017). Examining the construct validity of the MMPI-2-RF Interpersonal Functioning Scales using the Computerized Adaptive Test of Personality Disorder as a comparative framework. *Journal of Personality Assessment*, *99*, 416–423. doi: [10.1080/00223891.2016.1222394](https://doi.org/10.1080/00223891.2016.1222394)
- Goodwin, B.E., Sellbom, M., & Salekin, R. T. (2015). Elucidating the construct validity of the Antisocial Process Screening Device (APSD) in a sample of young adults. *Journal of Psychopathology and Behavioral Assessment*, *37*, 1–11. doi: [10.1007/s10862-014-9444-z](https://doi.org/10.1007/s10862-014-9444-z)
- Hoerger, M., Quirk, S. W., & Weed, N. C. (2011). Development and validation of the Delaying Gratification Inventory. *Psychological Assessment*, *23*, 725–738. doi: [10.1037/a0023286](https://doi.org/10.1037/a0023286)
- Hunter, H. K., Bolinsky, P. K., Novi, J. H., Hudak, D. V., James, A. V., Myers, K. R., & Schuder, K. M. (2014). Using the MMPI-2-RF to discriminate psychometrically identified schizotypic college students from a matched comparison sample. *Journal of Personality Assessment*, *96*, 596–603. doi: [10.1080/00223891.2014.922093](https://doi.org/10.1080/00223891.2014.922093)

- Ingram, P. B., Kelso, K. M., & McCord, D. M. (2011). Empirical correlates and expanded interpretation of the MMPI-2-RF Restructured Clinical Scale 3 (Cynicism). *Assessment, 18*, 95–101. doi: [10.1177/1073191110388147](https://doi.org/10.1177/1073191110388147)
- Johnson, A. K., Sellbom, M., & Glenn, A. L. (2018). Dimensional personality traits broadly and selectively associated with normative externalizing behavior. *Journal of Psychopathology and Behavioral Assessment*. doi: [10.1007/s10862-018-9665-7](https://doi.org/10.1007/s10862-018-9665-7)
- Kastner, R. M., & Sellbom, M. (2012). Hypersexuality in college students: The role of psychopathy. *Personality and Individual Differences, 53*, 644–649. doi: [10.1016/j.paid.2012.05.005](https://doi.org/10.1016/j.paid.2012.05.005)
- Kim, S., Goodman, G. M., Toruno, J. A., Sherry, A. R., & Kim, H. K. (2014). The cross-cultural validity of the MMPI-2-RF Higher-Order Scales in a sample of North Korean female refugees. *Assessment, 22*, 640–649. doi: [10.1177/1073191114548444](https://doi.org/10.1177/1073191114548444)
- Kim, S., Kim, H. K., & Lee, Narae. (2013). Psychological features of North Korean female refugees on the MMPI-2: A latent profile analysis. *Psychological Assessment 25*, 1091–1102. doi: [10.1037/a0033097](https://doi.org/10.1037/a0033097)
- Lange, R. T., Edmed, S. L., Sullivan, K. A., French, L. M., & Cooper, D. B. (2013). Utility of the Mild Brain Injury Atypical Symptoms Scale to detect symptom exaggeration. An analogue simulation study. *Journal of Clinical and Experimental Neuropsychology*. doi: [10.1080/13803395.2012.761677](https://doi.org/10.1080/13803395.2012.761677)
- Lee, T. T. C., & Forbey, J. D. (2010). MMPI-2 correlates of sexual preoccupation as measured by the Sexuality Scale in a college setting. *Sexual Addiction and Compulsivity, 17*, 219–235. doi: [10.1080/10720162.2010.500500](https://doi.org/10.1080/10720162.2010.500500)
- Lijewski, A., MacDonald, D. A., Paynard, C. M. (2013). Examination of the psychometric properties of the MMPI-2 Restructured Clinical (RC) Scales with a sample of public safety officer candidates. *The International Journal of Educational and Psychological Assessment, 13*, 1–12.
- McCord, D. M., Achee, M. C., Cannon, E. M., Harrop, T. M., & Poynter, D. M. (2017). Using the Research Domain Criteria framework to explore associations between MMPI-2-RF constructs and physiological variables assessed by eye-tracker technology. *Journal of Personality Assessment, 99*, 363–374. doi: [10.1080/00223891.2016.1228067](https://doi.org/10.1080/00223891.2016.1228067)
- Mendez, D. M., & MacDonald, D. A. (2012). Spirituality and the MMPI-2 Restructured Clinical Scales. *International Journal of Transpersonal Studies, 31*, 1–12.
- Miller, K. E., Koffel, E., Kramer, M. D., Erbes, C. R., Arbisi, P. A., & Polusny, M. A. (2018). At-home partner sleep functioning over the course of military deployment. *Journal of Family Psychology, 32*, 114–122. doi: [10.1037/fam0000262](https://doi.org/10.1037/fam0000262)
- Monaghan, C., Bizumic, B., & Sellbom, M. (in press). The role of Machiavellian views and tactics in psychopathology. *Personality and Individual Differences, 94*, 72–81. doi: [10.1016/j.paid.2016.01.002](https://doi.org/10.1016/j.paid.2016.01.002)
- Neal, T. M. S., & Sellbom, M. (2012). Examining the factor structure of the Hare Self-Report Psychopathy Scale. *Journal of Personality Assessment, 93*, 244–253. doi: [10.1080/00223891.2011.648294](https://doi.org/10.1080/00223891.2011.648294)
- Osberg, T. M., Haseley, E. N., & Kamas, M. M. (2008). The MMPI-2 Clinical Scales and Restructured Clinical (RC) Scales: Comparative psychometric properties and relative diagnostic efficiency in young adults. *Journal of Personality Assessment, 90*, 81–92. doi: [10.1080/00223890701693801](https://doi.org/10.1080/00223890701693801)
- Rinaldi, L., Locati, F., Parolin, L., & Girelli, L. (2017). Distancing the present self from the past and the future: Psychological distance in anxiety and depression. *The Quarterly Journal of Experimental Psychology, 70:7*, 1106–1113, doi: [10.1080/17470218.2016.1271443](https://doi.org/10.1080/17470218.2016.1271443)



Minnesota Multiphasic  
Personality Inventory-2  
Restructured Form®

## MMPI-2-RF® Bibliography continued

Fall 2018

- Schuder, K. M., Gooding, D. C., Matts, C. W., & Bolinskey, P. K. (2016). Further evidence of the MMPI-2-RF's ability to discriminate psychometrically identified schizotypic college students from a matched comparison sample. *Personality and Individual Differences, 94*, 107-112. doi: [10.1016/j.paid.2016.01.014](https://doi.org/10.1016/j.paid.2016.01.014)
- Sellbom, M. (in press). Elucidating the complex associations between psychopathy and post-traumatic stress disorder from the perspective of trait negative affectivity. *International Journal of Forensic Mental Health*.
- Sellbom, M., & Ben-Porath, Y. S. (2005). Mapping the MMPI-2 Restructured Clinical (RC) Scales onto normal personality traits: Evidence of construct validity. *Journal of Personality Assessment, 85*, 179-187. doi: [10.1207/s15327752jpa8502\\_10](https://doi.org/10.1207/s15327752jpa8502_10)
- Sellbom, M., Ben-Porath, Y. S., Lilienfeld, S. O., Patrick, C. J., & Graham, J. R. (2005). Assessing psychopathic personality traits with the MMPI-2. *Journal of Personality Assessment, 85*, 334-343. doi: [10.1207/s15327752jpa8503\\_10](https://doi.org/10.1207/s15327752jpa8503_10)
- Sellbom, M., & Smith, A. (2017). Assessment of DSM-5 Section II personality disorders with the MMPI-2-RF in a nonclinical sample. *Journal of Personality Assessment, 99*, 384-397. doi: [10.1080/00223891.2016.1242074](https://doi.org/10.1080/00223891.2016.1242074)
- Steenhaut, P., Demeyer, I., De Raedt, R., & Rossi, G. (2017). The role of personality in the assessment of subjective wellbeing and physiological emotional reactivity: A comparison between younger and older adults. *Assessment*. doi: [10.1177/1073191117719510](https://doi.org/10.1177/1073191117719510)